

Rabbit anti-Human GRP78/HSPA5 Polyclonal Antibody | anti-GRP78/HSPA5 antibody

Anti-GRP78/HSPA5 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258777	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry


PRODUCT SYNONYMS

GRP78/HSPA5; Polyclonal Antibody; Anti-GRP78/HSPA5 Antibody; Rabbit Polyclonal; heat shock 70kDa protein 5 (glucose-regulated protein; 78kDa); Anti-BIP Antibody; Anti-GRP78 Antibody; Anti-HEL-S-89n Antibody; Anti-MIF2 Antibody; anti-GRP78/HSPA5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human GRP78/HSPA5
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-GRP78/HSPA5 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to the N-terminus of the Human GRP78/HSPA5
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human GRP78/HSPA5, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human GRP78 in human prostatic carcinoma with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.




IHC (Immunohistochemistry)

Immunochemical staining of human GRP78 in human testis with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.




IHC (Immunohistochemistry)

Immunochemical staining of human GRP78 in mouse testis with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human GRP78 in rat testis with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IF (Immunofluorescence)

Immunofluorescence staining of HSPA5 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human HSPA5 polyclonal antibody (dilution ratio 1:100) at 4? overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/grp78-hspa5-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/258777>