

Rabbit anti-Human UACA Polyclonal Antibody | anti-UACA antibody

Anti-UACA Antibody, Rabbit Polyclonal

CATALOG # AAA258974	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

UACA; Polyclonal Antibody; Anti-UACA Antibody; Rabbit Polyclonal; uveal autoantigen with coiled-coil domains and ankyrin repeats; Anti-NUCLING Antibody; anti-UACA antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human UACA
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-UACA antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human UACA fragment
Preparation	Produced in rabbits immunized with E coli-derived Human UACA fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human UACA in human gallbladder with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human UACA in human kidney with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-UACA rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate
Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 162 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/uaca-antibody-immunohistochemistry-western-blot-elisa/258974>