

Rabbit anti-Human GTF2E2 Polyclonal Antibody | anti-GTF2E2 antibody

Anti-GTF2E2 Antibody, Rabbit Polyclonal


CATALOG # AAA258993	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

GTF2E2; Polyclonal Antibody; Anti-GTF2E2 Antibody; Rabbit Polyclonal; general transcription factor IIE subunit 2; Anti-FE Antibody; Anti-TF2E2 Antibody; Anti-TFIIIE-B Antibody; anti-GTF2E2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human GTF2E2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-GTF2E2 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E coli-derived Human GTF2E2 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human GTF2E2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IP (Immunoprecipitation)

GTF2E2 was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate 4 uL anti-GTF2E2 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-GTF2E2 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 33 kDa Observed band size :36 kDa



WB (Western Blot)

Anti-GTF2E2 rabbit polyclonal antibody at 1:500 dilution Lane A: K562 Whole Cell Lysate Lane B: Jurkat Whole Cell Lysate Lane C: 293T Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 33 kDa Observed band size: 35 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.