

Rabbit anti-Mouse CD47 Polyclonal Antibody | anti-CD47 antibody

Anti-CD47 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259144	Rabbit	Mouse	Immunoprecipitation, ELISA, Western Blot


PRODUCT SYNONYMS

CD47; Polyclonal Antibody; Anti-CD47 Antibody; Rabbit Polyclonal; CD47 molecule; Anti-9130415E20Rik Antibody; Anti-AA407862 Antibody; Anti-AI848868 Antibody; Anti-AW108519 Antibody; Anti-B430305P08Rik Antibody; Anti-IAP Antibody; Anti-Itgp Antibody; anti-CD47 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Mouse CD47
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-CD47 antibody	IP (Immunoprecipitation), ELISA, WB (Western Blot)
Immunogen	Recombinant Mouse CD47 Protein
Preparation	Produced in rabbits immunized with purified, recombinant Mouse CD47 (rM CD47). CD47 specific IgG was purified by Mouse CD47 affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



WB (Western Blot)

Anti-CD47 rabbit polyclonal antibody at 1:500 dilution Lane A: Mouse brain tissue lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 35 kDa



IP (Immunoprecipitation)

Mcd47 was immunoprecipitated using: Lane A: 0.5 mg Mouse Brain Whole Cell Lysate 4 uL anti-Mcd47 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-Mcd47 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 33 kDa Observed band size : 37 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-CD47 antibody	CD47 contains 1 Ig-like V-type (immunoglobulin-like) domain and is a receptor for the C-terminal cell binding domain of thrombospondin. It may play a role in membrane transport and signal transduction. CD47 is also a membrane protein, which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. It is very broadly distributed on normal adult tissues, as well as ovarian tumors, being especially abundant in some epithelia and the brain. CD47 may play a role in membrane transport and/or integrin dependent signal transduction. It may prevent premature elimination of red blood cells. It also may be involved in membrane permeability changes induced following virus infection. By acting as an adhesion receptor for THBS1 on platelets, CD47 plays a role in both cell adhesion and in the modulation of integrins. It also plays an important role in memory formation and synaptic plasticity in the hippocampus.
References	Brown EJ, et al. (2001) Integrin-associated protein (CD47) and its ligands. Trends Cell Biol. 11(3): 130-5. Oldenborg PA. (2004) Role of CD47 in erythroid cells and in autoimmunity. Leuk Lymphoma. 45(7): 1319-27. Kaczorowski DJ, et al. (2007) Targeting CD47: NO limit on therapeutic potential. Circ Res. 100(5): 602-3.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd47-antibody-immunoprecipitation-western-blot-elisa/259144>