

Rabbit anti-Rat EphA7 Polyclonal Antibody | anti-EphA7 antibody

Anti-EphA7 Antibody, Rabbit Polyclonal

CATALOG # AAA259147	HOST Rabbit	REACTIVITY Rat	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	--------------------------	--


PRODUCT SYNONYMS

EphA7; Polyclonal Antibody; Anti-EphA7 Antibody; Rabbit Polyclonal; Eph receptor A7; Anti-Ehk-3 Antibody; Anti-Ehk3 Antibody; anti-EphA7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Rat EphA7
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-EphA7 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Rat EphA7 Protein
Preparation	Produced in rabbits immunized with purified, recombinant Rat EphA7 (rR EphA7). EphA7 specific IgG was purified by Rat Cystatin C affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of Mouse Carbonic anhydrase II in Mouse brain with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of Mouse Carbonic anhydrase II in Mouse kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-EphA7 antibody	Ephrin type-A receptor 7, also known as EphA7, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity6. Eph receptor-mediated signaling, which is triggered by ephrins7, probably modifies the properties of synapses during synaptic activation and remodeling. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). Ligand-mediated activation of Ephs induces various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer. Down-regulation of EphA7 secondary to hypermethylation has been reported in colorectal cancer. The expression of EphA7 was reduced in all tested gastric cancer cell lines; however, there is marked variability in expression among gastric carcinoma specimens. EphA7 may have roles in the pathogenesis and development of gastric carcinomas.
References	Rashid T, et al. (2005) Opposing gradients of ephrin-As and EphA7 in the superior colliculus are essential for topographic mapping in the mammalian visual system. Neuron. 47(1): 57-69. Wang J, et al. (2007) Differential expression of EphA7 receptor tyrosine kinase in gastric carcinoma. Hum Pathol. 38(11): 1649-56. Rogers JH, et al. (1999) Distribution of the receptor EphA7 and its ligands in development of the mouse nervous system. Brain Res Mol Brain Res. 74(1-2): 225-30.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/epha7-antibody-immunohistochemistry-elisa/259147>