

Rabbit anti-Human EIF2B4 Polyclonal Antibody | anti-EIF2B4 antibody

Anti-EIF2B4 Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259530	Rabbit	Human	Immunocytochemistry, Western Blot

PRODUCT SYNONYMS


EIF2B4; Polyclonal Antibody; Anti-EIF2B4 Antibody; Rabbit Polyclonal; EIF2B4 Antibody; Rabbit PAb; Antigen Affinity Purified; eukaryotic translation initiation factor 2B; subunit 4 delta; 67kDa; Anti-DKFZP586J0119 Antibody; Anti-EIF-2B Antibody; Anti-EIF2B Antibody; Anti-EIF2BD Antibody; Anti-EIF2Bdelta Antibody; Anti-VWM4 Antibody; anti-EIF2B4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human EIF2B4
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-EIF2B4 antibody	ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E coli-derived Human EIF2B4 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human EIF2B4 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2 degree C-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

**IF (Immunofluorescence)**

Immunofluorescence staining of EIF2B4—FP09827 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human EIF2B4—FP09827 polyclonal antibody (dilution ratio 1:100) at 4 degree C overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.

**WB (Western Blot)**

Anti-EIF2B4 rabbit polyclonal antibody at 1:500 dilution Lane A: K562 Whole Cell Lysate Lysates/proteins at 30ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 57 kDa Observed band size: 57 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/eif2b4-antibody-immunocytochemistry-western-blot-elisa/259530>