

Mouse anti-Human CD16 Monoclonal Antibody | anti-FCGR3B antibody

Anti-CD16 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260049	Mouse	Human	Flow Cytometry, Functional Assay, Immunofluorescence


PRODUCT SYNONYMS

CD16; Monoclonal Antibody; Anti-CD16 Antibody; CD16B; FCG3; FCGR3; IGFR3; Low affinity immunoglobulin gamma Fc region receptor III-B; Fc-gamma RIII-beta; Fc-gamma RIII; Fc-gamma RIIIB; FcRIII; FcRIIIB; FcR-10; IgG Fc receptor III-1; CD antigen CD16b; anti-FCGR3B antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Specificity	Recognises human CD16.
Purity/Purification	The antibody was purified by affinity chromatography.
Form/Format	Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Sequence Length	233
Applicable Applications for anti-FCGR3B antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence)
Directions for Use	1. Take 100 ul peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add appropriate amount of antibody to the bottom of flow tube mixing with the whole blood; incubate for 30 minutes at room temperature. 3. Add 2 ml RBC lysis buffer; incubate for 10 minutes after mixing; dissolve red blood cells. 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes; discard the supernatant. 5. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 6. Add appropriate amount of fluorescent-labeled anti-mouse IgGs and incubate for 20 minutes away from light at room temperature. 7. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 8. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES



FCM/FACS (Flow Cytometry)
Flow cytometric analysis of human peripheral blood lymphocytes using Anti-CD16 Antibody, followed by anti-mouse IgG PE.

ADDITIONAL INFORMATION

Related Product Information for anti-FCGR3B antibody	Mouse monoclonal antibody to CD16
--	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	132814489
NCBI GeneID	2215
NCBI Accession #	NP_000561.3
NCBI GenBank Nucleotide #	NM_000570.4
UniProt Accession #	O75015
NCBI Official Full Name	low affinity immunoglobulin gamma Fc region receptor III-B isoform 2
NCBI Official Synonym Full Names	Fc fragment of IgG, low affinity IIIb, receptor (CD16b)
NCBI Official Symbol	FCGR3B
NCBI Official Synonym Symbols	CD16; FCG3; CD16b; FCGR3; FCR-10; FCRIII; FCRIIIb
NCBI Protein Information	low affinity immunoglobulin gamma Fc region receptor III-B
UniProt Protein Name	Low affinity immunoglobulin gamma Fc region receptor III-B
UniProt Gene Name	FCGR3B
UniProt Synonym Gene Names	CD16B; FCG3; FCGR3; IGFR3; Fc-gamma RIII; Fc-gamma RIIIb; FcRIII; FcRIIIb
UniProt Entry Name	FCG3B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd16-antibody-flow-cytometry-immunofluorescence-elisa/260049>