

Rabbit BMX Polyclonal Antibody | anti-BMX antibody

Anti-BMX Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260554	Rabbit	Human, Mouse, Rat, Bovine, Dog, Pig	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

BMX; Polyclonal Antibody; Anti-BMX Antibody; Cytoplasmic tyrosine-protein kinase BMX; Bone marrow tyrosine kinase gene in chromosome X protein; Epithelial and endothelial tyrosine kinase; ETK; NTK38; anti-BMX antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of BMX protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	675
Applicable Applications for anti-BMX antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human BMX. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of BMX staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of BMX staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of BMX expression in A549 (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-BMX antibody

Rabbit polyclonal antibody to BMX

NCBI AND UNIPROT INFORMATION

NCBI GI #	1705489
NCBI GeneID	660
UniProt Accession #	P51813; O60564; Q12871; A6NIH9
NCBI Official Full Name	Cytoplasmic tyrosine-protein kinase BMX
NCBI Official Synonym Full Names	BMX non-receptor tyrosine kinase
NCBI Official Symbol	BMX
NCBI Official Synonym Symbols	ETK; PSCTK2; PSCTK3
NCBI Protein Information	cytoplasmic tyrosine-protein kinase BMX
UniProt Protein Name	Cytoplasmic tyrosine-protein kinase BMX
UniProt Gene Name	BMX
UniProt Synonym Gene Names	ETK
UniProt Entry Name	BMX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/bmx-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/260554>