

Mouse IL-12 p70 ELISA Kit | IL12A elisa kit

Mouse IL-12 p70 ELISA Kit

CATALOG # AAA261652	REACTIVITY Mouse	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

IL-12 p70; Mouse IL-12 p70 ELISA Kit; NKSF1; Interleukin-12 subunit alpha; IL-12A; Cytotoxic lymphocyte maturation factor 35 kDa subunit; CLMF p35; IL-12 subunit p35; NK cell stimulatory factor chain 1; NKSF2; Interleukin-12 subunit beta; IL-12B; Cytotoxic lymphocyte maturation factor 40 kDa subunit; CLMF p40; IL-12 subunit p40; NK cell stimulatory factor chain 2; IL12A elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	The Mouse IL-12 p70 ELISA Kit allows for the detection and quantification of endogenous levels of natural and/or recombinant Mouse IL-12 p70 proteins within the range of 15.6 pg/ml - 1000 pg/ml.
Sequence Length	215
Samples	Cell Culture Supernatants, Serum, Plasma, Tissue Homogenates.
Sensitivity	7 pg/ml.
Applicable Applications for IL12A elisa kit	ELISA (Sandwich ELISA)
Preparation and Storage	Shipped and store at 4 degree C for 6 months, store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL12A elisa kit	Principle of the Assay The Mouse IL-12 p70 ELISA (Enzyme-Linked Immunosorbent Assay) kit is an in vitro enzyme-linked immunosorbent assay for the quantitative measurement of Mouse IL-12 p70 in Cell Culture Supernatants, Serum, Plasma, Tissue Homogenates. This assay employs an antibody specific for Mouse IL-12 p70 coated on a 96-well plate. Standards and samples are pipetted into the wells and IL-12 p70 present in a sample is bound to the wells by the immobilized antibody. The wells are washed and biotinylated anti-Mouse IL-12 p70 antibody is added. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin is pipetted to the wells. The wells are again washed, a TMB substrate solution is added to the wells and color develops in proportion to the amount of IL-12 p70 bound. The Stop Solution changes the color from blue to yellow, and the intensity of the color is measured at 450 nm. Background/Introduction: Interleukin (IL)-12 is a 70-KDa cytokine comprised of two disulfide-linked proteins (p35 and p40) and is essential for the initiation of effective immune response. And the IL-12 p70 is a heterodimer of p35 and p40 subunits; it is an important cytokine secreted by antigen-presenting cells in response to antigenic stimulation. Gene expression analysis of the IL-12 cytokine family subunits revealed that both strains induced high levels of p40 (protein chain communal to IL-12 p70 and IL-23) as well as p19, a subunit of IL-23. Conversely only ACT- 18HS19 infection induced consistent transcription of IL-12 p35, a subunit of IL-12 p70.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	1170458
NCBI GeneID	16159
UniProt Accession #	P43431; P43432
Molecular Weight	24,179 Da
NCBI Official Full Name	Interleukin-12 subunit alpha
NCBI Official Synonym Full Names	interleukin 12a
NCBI Official Symbol	IL12a
NCBI Official Synonym Symbols	p35; LI12a; IL-12a; IL-12p35
NCBI Protein Information	interleukin-12 subunit alpha
UniProt Protein Name	Interleukin-12 subunit alpha
UniProt Gene Name	IL12a
UniProt Synonym Gene Names	IL-12A; CLMF p35
UniProt Entry Name	IL12A_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.