

Human CD274 ELISA Kit | CD274 elisa kit

Human CD274 ELISA Kit

CATALOG #
AAA263026

REACTIVITY
Human


PRODUCT SYNONYMS

CD274; Human CD274 ELISA Kit; B7H1; PDCD1L1; PDCD1LG1; PDL1; Programmed cell death 1 ligand 1; PD-L1; PDCD1 ligand 1; Programmed death ligand 1; B7 homolog 1; B7-H1; CD274 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	The Human CD274 ELISA Kit allows for the detection and quantification of endogenous levels of natural and/or recombinant Human CD274 proteins within the range of 62.5 pg/ml - 4000 pg/ml.
Sequence Length	176
Samples	Cell Culture Supernates, Serum, Plasma
Sensitivity	<12 pg/ml
Preparation and Storage	Shipped and store at 4 degree C for 6 months, store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD274 elisa kit	Principle of the Assay: The Human CD274 ELISA (Enzyme-Linked Immunosorbent Assay) kit is an in vitro enzyme-linked immunosorbent assay for the quantitative measurement of Human CD274 in Cell Culture Supernates, Serum, Plasma. This assay employs an antibody specific for Human CD274 coated on a 96-well plate. Standards and samples are pipetted into the wells and CD274 present in a sample is bound to the wells by the immobilized antibody. The wells are washed and biotinylated anti-Human CD274 antibody is added. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin is pipetted to the wells. The wells are again washed, a TMB substrate solution is added to the wells and color develops in proportion to the amount of CD274 bound. The Stop Solution changes the color from blue to yellow, and the intensity of the color is measured at 450 nm. Background: Programmed death-ligand 1 (PD-L1), also known as cluster of differentiation 274 (CD274) or B7 homolog 1 (B7-H1), is a protein that in humans is encoded by the CD274 gene. PD-L1 is a 40kDa type 1 transmembrane protein that has been speculated to play a major role in suppressing the immune system during particular events such as pregnancy, tissue allografts, autoimmune disease and other disease states such as hepatitis. Normally the immune system reacts to foreign antigens where there is some accumulation in the lymph nodes or spleen which triggers a proliferation of antigen-specific CD8+ T cell. The formation of PD-1 receptor / PD-L1 or B7.1 receptor /PD-L1 ligand complex transmits an inhibitory signal which reduces the proliferation of these CD8+ T cells at the lymph nodes and supplementary to that PD-1 is also able to control the accumulation of foreign antigen specific T cells in the lymph nodes through apoptosis which is further mediated by a lower regulation of the gene Bcl-2.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	390979639
NCBI GeneID	29126
NCBI Accession #	NP_001254635.1
NCBI GenBank Nucleotide #	NM_001267706.1
UniProt Accession #	Q9NZQ7; Q14CJ2; Q2V8D5; Q66RK1; Q6WEX4; Q9NUZ5; B2RBA2; B4DU27
Molecular Weight	20,454 Da
NCBI Official Full Name	programmed cell death 1 ligand 1 isoform b
NCBI Official Synonym Full Names	CD274 molecule
NCBI Official Symbol	CD274
NCBI Official Synonym Symbols	B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1
NCBI Protein Information	programmed cell death 1 ligand 1
UniProt Protein Name	Programmed cell death 1 ligand 1
UniProt Gene Name	CD274
UniProt Synonym Gene Names	B7H1; PDCD1L1; PDCD1LG1; PDL1; PD-L1; PDCD1 ligand 1; Programmed death ligand 1; B7-H1
UniProt Entry Name	PD1L1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.