

Human Epiregulin ELISA Kit | EREG elisa kit

Human Epiregulin ELISA Kit

CATALOG #
AAA263041

REACTIVITY
Human


PRODUCT SYNONYMS

Epiregulin; Human Epiregulin ELISA Kit; Proepiregulin; EREG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	The Human Epiregulin ELISA Kit allows for the detection and quantification of endogenous levels of natural and/or recombinant Human Epiregulin proteins within the range of 156 pg/ml - 10000 pg/ml.
Sequence Length	169
Samples	Cell Culture Supernates, Serum, Plasma
Sensitivity	<10 pg/ml
Preparation and Storage	Shipped and store at 4 degree C for 6 months, store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for EREG elisa kit

Principle of the Assay: The Human Epiregulin ELISA (Enzyme-Linked Immunosorbent Assay) kit is an in vitro enzyme-linked immunosorbent assay for the quantitative measurement of Human Epiregulin in Cell Culture Supernates, Serum, Plasma. This assay employs an antibody specific for Human Epiregulin coated on a 96-well plate. Standards and samples are pipetted into the wells and Epiregulin present in a sample is bound to the wells by the immobilized antibody. The wells are washed and biotinylated anti-Human Epiregulin antibody is added. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin is pipetted to the wells. The wells are again washed, a TMB substrate solution is added to the wells and color develops in proportion to the amount of Epiregulin bound. The Stop Solution changes the color from blue to yellow, and the intensity of the color is measured at 450 nm.

Background: Epiregulin (EPR) is a protein that in humans is encoded by the EREG gene. It is a member of the epidermal growth factor family. Epiregulin can function as a ligand of epidermal growth factor receptor (EGFR), as well as a ligand of most members of the ERBB (v-erb-b2 oncogene homolog) family of tyrosine-kinase receptors. The secondary structure at the C-terminus epiregulin is different from other epidermal growth factor family ligands because of the lack of hydrogen bonds. The structural difference at the C-terminus may provide an explanation for the reduced binding affinity of epiregulin to the ERBB receptors.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557567
NCBI GeneID	2069
NCBI Accession #	NP_001423.1
NCBI GenBank Nucleotide #	NM_001432.2
UniProt Accession #	O14944; Q6FH69; B2RC66
Molecular Weight	19,044 Da
NCBI Official Full Name	proepiregulin preproprotein
NCBI Official Synonym Full Names	epiregulin
NCBI Official Symbol	EREG
NCBI Official Synonym Symbols	ER; Ep; EPR
NCBI Protein Information	proepiregulin
UniProt Protein Name	Proepiregulin
UniProt Gene Name	EREG
UniProt Synonym Gene Names	EPR
UniProt Entry Name	EREG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/ereg-human-elisa-kits/epiregulin/263041>