

General NADH Dehydrogenase Assay Kit | NADH assay kit

NADH Dehydrogenase Microplate Assay Kit

CATALOG # AAA264766	REACTIVITY General	APPLICATIONS Functional Assay
-------------------------------	------------------------------	---


PRODUCT SYNONYMS

NADH Dehydrogenase; NADH Dehydrogenase Microplate Assay Kit; EC 1.6.5.3; ubiquinone reductase (ambiguous); type 1 dehydrogenase; complex 1 dehydrogenase; coenzyme Q reductase (ambiguous); complex I (electron transport chain); complex I (mitochondrial electron transport); complex I (NADH:Q1 oxidoreductase); dihydronicotinamide adenine dinucleotide-coenzyme Q reductase (ambiguous); DPNH-coenzyme Q reductase (ambiguous); DPNH-ubiquinone reductase (ambiguous); mitochondrial electron transport complex 1; mitochondrial electron transport complex I; NADH coenzyme Q1 reductase; NADH-coenzyme Q oxidoreductase (ambiguous); NADH-coenzyme Q reductase (ambiguous); NADH-CoQ oxidoreductase (ambiguous); NADH-CoQ reductase (ambiguous); NADH-ubiquinone reductase (ambiguous); NADH-ubiquinone oxidoreductase (ambiguous); NADH-ubiquinone-1 reductase; reduced nicotinamide adenine dinucleotide-coenzyme Q reductase (ambiguous); NADH:ubiquinone oxidoreductase complex; NADH-Q6 oxidoreductase (ambiguous); electron transfer complex I; NADH2 dehydrogenase (ubiquinone); NADH assay kit

PRODUCT SPECIFICATIONS

Reactivity	General
Samples	Tissue extracts, Cell lysate, Cell culture media and Other biological fluids Samples
Detection Range	4 umol/L - 400 umol/L
Applicable Applications for NADH assay kit	FA (Functional Assay)
Preparation and Storage	Shipped and store at 4 degree C for 6 months.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NADH assay kit

Background: NADH dehydrogenase (EC 1.6.5.3) (also referred to as NADH:ubiquinone oxidoreductase or, especially in the context of the human protein, Complex I) is an enzyme of the respiratory chains of myriad organisms from bacteria to humans that falls under the H⁺ or Na⁺-translocating NADH Dehydrogenase (NDH) Family, a member of the Na⁺ transporting Mrp superfamily. It catalyzes the transfer of electrons from NADH to coenzyme Q10 (CoQ10) and, in eukaryotes, it is located in the inner mitochondrial membrane. NADH:ubiquinone oxidoreductases type I of bacteria and of eukaryotic mitochondria and chloroplasts couple electron transfer to the electrogenic transport of protons or Na⁺. It is one of the "entry enzymes" of cellular respiration or oxidative phosphorylation in the mitochondria. The assay is used to determine NADH Dehydrogenase. The enzyme catalysed reaction products is NAD⁺. The reaction velocity is measured as a decrease in A340 resulting from the oxidation of NADH.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/general-assay-kits/nadh-dehydrogenase/264766>