

Mouse anti-Human CD23 Monoclonal Antibody | anti-FCER2 antibody

Anti-CD23 Antibody

CATALOG # AAA264821	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

CD23; Monoclonal Antibody; Anti-CD23 Antibody; CD23A; CLEC4J; FCE2; IGEBF; Low affinity immunoglobulin epsilon Fc receptor; BLAST-2; C-type lectin domain family 4 member J; Fc-epsilon-RII; Immunoglobulin E-binding factor; Lymphocyte IgE receptor; anti-FCER2 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1
Specificity	Recognizes endogenous levels of CD23 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for anti-FCER2 antibody	IHC (Immunohistochemistry)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human CD23. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunohistochemical analysis of CD23 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD23 staining in human follicular lymphoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-FCER2 antibody	Mouse monoclonal antibody to CD23
---	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	119862
NCBI GeneID	2208
UniProt Accession #	P06734
Molecular Weight	321
NCBI Official Full Name	Low affinity immunoglobulin epsilon Fc receptor
NCBI Official Synonym Full Names	Fc fragment of IgE, low affinity II, receptor for (CD23)
NCBI Official Symbol	FCER2
NCBI Official Synonym Symbols	CD23; FCE2; CD23A; IGEBF; CLEC4J; BLAST-2
NCBI Protein Information	low affinity immunoglobulin epsilon Fc receptor; CD23 antigen; fc-epsilon-RII; lymphocyte IgE receptor; immunoglobulin E-binding factor; C-type lectin domain family 4, member J
UniProt Protein Name	Low affinity immunoglobulin epsilon Fc receptor
UniProt Gene Name	FCER2
UniProt Synonym Gene Names	CD23A; CLEC4J; FCE2; IGEBF
UniProt Entry Name	FCER2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd23-antibody-immunohistochemistry-elisa/264821>