

Rabbit CD4 Polyclonal Antibody | anti-CD4 antibody

Anti-CD4 (Phospho-S433) Antibody

CATALOG # AAA265046	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Dog, Monkey	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	---	---


PRODUCT SYNONYMS

CD4; Polyclonal Antibody; Anti-CD4 (Phospho-S433) Antibody; anti-CD4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Dog, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of CD4 protein only when phosphorylated at S433.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-CD4 antibody	IC (Immunochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding S433 of human CD4 protein. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




ELISA

Direct ELISA antibody dose-response curve using Anti-CD4 (Phospho-S433) Antibody. Antigen (Phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.



IF (Immunofluorescence)

Immunofluorescent analysis of CD4 (Phospho-S433) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



WB (Western Blot)

Western blot analysis of CD4 (Phospho-S433) expression in mouse spleen (A), rat spleen (B) whole cell lysates. (Predicted band size: 51 kD; Observed band size: 56 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-CD4 antibody	Rabbit polyclonal antibody to CD4 (Phospho-S433)
Product Categories/Family for anti-CD4 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	10835167
NCBI GeneID	920
NCBI Accession #	NP_000607.1
NCBI GenBank Nucleotide #	NM_000616.4
UniProt Accession #	P01730 ; P06332 ; P05540
Molecular Weight	51,111 Da
NCBI Official Full Name	T-cell surface glycoprotein CD4 isoform 1
NCBI Official Synonym Full Names	CD4 molecule
NCBI Official Symbol	CD4
NCBI Official Synonym Symbols	CD4mut
NCBI Protein Information	T-cell surface glycoprotein CD4; CD4 receptor; CD4 antigen (p55); T-cell surface antigen T4/Leu-3
UniProt Protein Name	T-cell surface glycoprotein CD4
UniProt Gene Name	CD4
UniProt Entry Name	CD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cd4-antibody-immunochemistry-immunofluorescence-western-blot-elisa/265046>