

Rabbit MUC2 Polyclonal Antibody | anti-MUC2 antibody

Anti-MUC2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265220	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MUC2; Polyclonal Antibody; Anti-MUC2 Antibody; anti-MUC2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MUC2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-MUC2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MUC2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MUC2 staining in mouse colon. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MUC2 staining in human colon formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MUC2 expression in mouse small intestine (A), rat small intestine (B) whole cell lysates. (Predicted band size: 540 kD; Observed band size: 110 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-MUC2 antibody	Rabbit polyclonal antibody to MUC2
Product Categories/Family for anti-MUC2 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	116284392
NCBI GeneID	4583
NCBI Accession #	NP_002448.2
NCBI GenBank Nucleotide #	NM_002457.2
UniProt Accession #	Q02817 ; Q80Z19 ; Q62635
Molecular Weight	540,300 Da
NCBI Official Full Name	mucin-2
NCBI Official Synonym Full Names	mucin 2, oligomeric mucus/gel-forming
NCBI Official Symbol	MUC2
NCBI Official Synonym Symbols	MLP; SMUC; MUC-2
NCBI Protein Information	mucin-2; intestinal mucin-2; mucin-like protein; mucin 2, intestinal/tracheal
UniProt Protein Name	Mucin-2
UniProt Gene Name	MUC2
UniProt Synonym Gene Names	SMUC; MUC-2
UniProt Entry Name	MUC2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.