

Mouse AKT2 Monoclonal Antibody | anti-AKT2 antibody

AKT2 (v-akt Murine Thymoma Viral Oncogene Homolog 2, PKBB, PKBBETA, PRKBB, RAC-BETA) (HRP)


CATALOG #	HOST	APPLICATIONS
AAA26545	Mouse	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


AKT2; Monoclonal Antibody; AKT2 (v-akt Murine Thymoma Viral Oncogene Homolog 2; PKBB; PKBBETA; PRKBB; RAC-BETA) (HRP); v-akt Murine Thymoma Viral Oncogene Homolog 2; RAC-BETA; anti-AKT2 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,k
Clone Number	X1
Specificity	Recognizes AKT2.
Purity/Purification	Purified
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with horseradish peroxidase (HRP).
Sequence Length	481
Applicable Applications for anti-AKT2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Application Notes	Applications are based on unconjugated antibody.
Immunogen	AKT2 (AAA58364, 100aa-189aa) partial recombinant protein with GST tag. MW of the GST tag alone is 26kD.
Immunogen Sequence	MRAIQMVANSLKQRAPGEDPMDYKCGSPSDSSTTEEMEVAVSKARAKVTMNDFDYLKLLGKGTFGKVILVREKATGRYYAMKILRKEVII
Conjugate	HRP
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and store at -20 degree C. Aliquots are stable at -20 degree C for 12 months after receipt. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Note: Sodium azide is a potent inhibitor of peroxidase and should not be added to HRP conjugates. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.




WB (Western Blot)
AKT2 monoclonal antibody (M06), clone X1. Western Blot analysis of AKT2 expression in PC-12.




WB (Western Blot)
AKT2 monoclonal antibody (M06), clone X1. Western Blot analysis of AKT2 expression in NIH/3T3.




IHC (Immunohistochemistry)
Immunoperoxidase of monoclonal antibody to AKT2 on formalin-fixed paraffin-embedded human small Intestine. [antibody concentration 3 ug/ml]



IHC (Immunohistochemistry)
Immunoperoxidase of monoclonal antibody to AKT2 on formalin-fixed paraffin-embedded human small Intestine. [antibody concentration 3 ug/ml]



Application Data
Detection limit for recombinant GST tagged AKT2 is approximately 1ng/ml as a capture antibody.



WB (Western Blot)
AKT2 monoclonal antibody (M06), clone X1 Western Blot analysis of AKT2 expression in Jurkat (Cat # L017V1).

ADDITIONAL INFORMATION

Product Categories/Family for anti-AKT2 antibody	Antibodies; Protein Kinases
--	-----------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	178326
NCBI GeneID	208

NCBI Official Full Name	protein serine/threonine kinase
NCBI Official Synonym Full Names	AKT serine/threonine kinase 2
NCBI Official Symbol	AKT2
NCBI Official Synonym Symbols	PKBB; PRKBB; HIHGHH; PKBBETA; RAC-BETA
NCBI Protein Information	RAC-beta serine/threonine-protein kinase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/akt2-antibody-clone-x1-immunofluorescence-immunohistochemistry-western-blot-elisa/26545>