

Rabbit MMP2 Polyclonal Antibody | anti-MMP2 antibody

Anti-MMP2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA269597	Rabbit	Human, Mouse, Rat, Bovine, Dog, Pig	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MMP2; Polyclonal Antibody; Anti-MMP2 Antibody; CLG4A; 72 kDa type IV collagenase; 72 kDa gelatinase; Gelatinase A; Matrix metalloproteinase-2; MMP-2; TBE-1; anti-MMP2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-MMP2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MMP2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MMP2 staining in COLO205 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MMP2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MMP2 expression in HuvEc (A), NRK (B), MCF7 (C), HT1080 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MMP2 antibody	Rabbit polyclonal antibody to MMP2
Product Categories/Family for anti-MMP2 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	189217853
NCBI GeneID	4313
NCBI Accession #	NP_001121363.1
NCBI GenBank Nucleotide #	NM_001127891.1
UniProt Accession #	P08253; Q9UCJ8; B2R6U1; B4DWH3; E9PE45
Molecular Weight	73,882 Da
NCBI Official Full Name	72 kDa type IV collagenase isoform b
NCBI Official Synonym Full Names	matrix metalloproteinase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase)
NCBI Official Symbol	MMP2
NCBI Official Synonym Symbols	CLG4; MONA; CLG4A; TBE-1; MMP-II
NCBI Protein Information	72 kDa type IV collagenase; MMP-2; gelatinase A; 72 kDa gelatinase; collagenase type IV-A; neutrophil gelatinase; matrix metalloproteinase-2; matrix metalloproteinase-II
UniProt Protein Name	72 kDa type IV collagenase
UniProt Gene Name	MMP2
UniProt Synonym Gene Names	CLG4A; MMP-2
UniProt Entry Name	MMP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mmp2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/269597>