

Rabbit ASC Polyclonal Antibody | anti-ASC antibody

Anti-ASC Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA269765	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ASC; Polyclonal Antibody; Anti-ASC Antibody; CARD5; TMS1; Apoptosis-associated speck-like protein containing a CARD; hASC; Caspase recruitment domain-containing protein 5; PYD and CARD domain-containing protein; Target of methylation-induced silencing 1; anti-ASC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-ASC antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ASC. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ASC staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of ASC staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ASC expression in HEK293T (A), Raw264.7 (B), mouse brain (C), rat brain (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ASC antibody	Rabbit polyclonal antibody to ASC
Product Categories/Family for anti-ASC antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	18203507
NCBI GeneID	29108
UniProt Accession #	Q9ULZ3; Q96D12; Q9BSZ5; Q9HBD0; Q9NXJ8
Molecular Weight	195
NCBI Official Full Name	Apoptosis-associated speck-like protein containing a CARD
NCBI Official Synonym Full Names	PYD and CARD domain containing
NCBI Official Symbol	PYCARD
NCBI Official Synonym Symbols	ASC; TMS; TMS1; CARD5; TMS-1
NCBI Protein Information	apoptosis-associated speck-like protein containing a CARD; target of methylation-induced silencing 1; caspase recruitment domain-containing protein 5
UniProt Protein Name	Apoptosis-associated speck-like protein containing a CARD
UniProt Gene Name	PYCARD
UniProt Synonym Gene Names	ASC; CARD5; TMS1; hASC
UniProt Entry Name	ASC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/asc-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/269765>