

Mouse PDGFRalpha Monoclonal Antibody | anti-PDGFRalpha antibody

PDGFRalpha (5D1) Mouse mAb

CATALOG # AAA276903	HOST Mouse	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	--	---

PRODUCT SYNONYMS

PDGFRalpha; Monoclonal Antibody; PDGFRalpha (5D1) Mouse mAb; anti-PDGFRalpha antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1
Form/Format	PBS with 0.02% sodium azide and 50% glycerol pH7.4.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-PDGFRalpha antibody	IHC (Immunohistochemistry)
Species	Human
Immunogen	Synthetic Peptide
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human Skin Tissue using PDGFR a(EM1336)Mouse mAb diluted at 1:200



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Rat Skin Tissue using PDGFR a (EM1336) Mouse mAb diluted at 1:200.

ADDITIONAL INFORMATION

Related Product Information for anti-PDGFRalpha antibody	Platelet derived growth factor (PDGF) family proteins exist as several disulphide-bonded, dimeric isoforms (PDGF AA, PDGF AB, PDGF BB, PDGF CC, and PDGF DD) that bind in a specific pattern to two closely related receptor tyrosine kinases, PDGF receptor alpha (PDGFRalpha) and PDGF receptor beta (PDGFRbeta). PDGFRalpha and PDGFRbeta can each form heterodimers with EGFR, which is also activated by PDGF. Various cells differ in the total number of receptors present and in the receptor subunit composition, which may account for responsive differences among cell types to PDGF binding. Ligand binding induces receptor dimerization and autophosphorylation, followed by binding and activation of cytoplasmic SH2 domain-containing signal transduction molecules, such as GRB2, Src, GAP, PI3 kinase, PLCgamma, and NCK. A number of different signaling pathways are initiated by activated PDGF receptors and lead to control of cell growth, actin reorganization, migration, and differentiation.
Product Categories/Family for anti-PDGFRalpha antibody	Membrane

NCBI AND UNIPROT INFORMATION

NCBI GI #	5453870
NCBI GeneID	5156
NCBI Accession #	NP_006197.1
NCBI GenBank Nucleotide #	NM_006206.4
UniProt Accession #	P16234; Q6P4H5; Q96KZ7; Q9UD28; B2RE69; E9PBH0
Molecular Weight	122,670 Da
NCBI Official Full Name	platelet-derived growth factor receptor alpha
NCBI Official Synonym Full Names	platelet-derived growth factor receptor, alpha polypeptide
NCBI Official Symbol	PDGFRA
NCBI Official Synonym Symbols	CD140A; PDGFR2; PDGFR-2; RHEPDGFRA
NCBI Protein Information	platelet-derived growth factor receptor alpha; PDGF-R-alpha; CD140a antigen; PDGFRA/BCR fusion; CD140 antigen-like family member A; platelet-derived growth factor receptor 2; alpha-type platelet-derived growth factor receptor; rearranged-in-hypereosinophi
UniProt Protein Name	Platelet-derived growth factor receptor alpha
UniProt Gene Name	PDGFRA
UniProt Synonym Gene Names	PDGFR2; RHEPDGFRA; PDGF-R-alpha; PDGFR-alpha; PDGFR-2
UniProt Entry Name	PGFRA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/pdgfralpha-antibody-immunohistochemistry-elisa/276903>