

Semaphorin-4D (SEMA4D) Active Protein | SEMA4D active protein

Recombinant Human Semaphorin-4D (SEMA4D), partial (Active)

CATALOG #
AAA278920

HOST
Mammalian Cell


PRODUCT SYNONYMS

Semaphorin-4D (SEMA4D); Recombinant Human Semaphorin-4D (SEMA4D); partial (Active); SEMA4D active protein


PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	22-734aa, Partial
Sequence	MAFAPIPRIT WEHREVHLVQ FHEPDIYNYS ALLLSEDKDT LYIGAREAVF AVNALNISEK QHEVYWKVSE DKKAKCAEKG KSKQTECLNY IRVLQPLSAT SLYVCGTNAF QPACDHLNLT SFKFLGKNED GKGRCPFDPA HSYTSVMVDG ELYSGTSYNF LGSEPIISRN SSHSPLRTEY AIPWLNPSF VFADVIRKSP DSPDGEDDRV YFFTEVSVE YEFVFRVLIP RIARVCKGDQ GGLRTLQKKW TSFLKARLIC SRPDSGLVFN VLRDVFVLR PGLKVPVFYA LFTPQLNNVG LSAVCAYNLS TAEVFSHGK YMQSTTVEQS HTKWVRYNGP VPKPRPGACI DSEARAANYT SSLNLPDKTL QFVKDHPLMD DSVTPIDNRP RLIKKDVNYT QIVVDRTQAL DGTYYDVMFV STDRGALHKA ISLEHAVHII EETQLFQDFE PVQTLTLLSSK KGNRFVYAGS NSGVVQAPLA FCGKHGTCED CVLARDPYCA WSPPTATCVA LHQTESPSRG LIQEMSGDAS VCPDKSKGSY RQHFFKHGGT AELKCSQKSN LARVFWKFQN GVLKAESPKY GLMGRKNLLI FNLSEGDGV YQCLSEERVK NKTVFQVVAK HVLEVKVVPK PVVAPTLVV QTEGSRIATK VLVASTQGSS PPTPAVQATS SGAILPPKP APTGTSCPEK IVINTVPQLH SEKTMYLKSS DNR
Species	Homo sapiens (Human)
Endotoxin	Less than 1.0EU/ug as determined by LAL method.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized human SEMA4D at 2ug/mL can bind biotinylated Pepinemab, the EC50 is 1.005-1.776ng/mL.
Preparation and Storage	Store at -20 degree C. For extended storage, store at -20 or -80 degree C. Repeated freezing and thawing is not recommended. Store working aliquots at 4 degree C for up to one week.

IMAGES



Bioactivity



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Product Categories/Family for SEMA4D active protein	Immunology
---	------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	214010218
NCBI GeneID	10507
NCBI Accession #	NP_006369.3
NCBI GenBank Nucleotide #	NM_006378.3
UniProt Accession #	Q92854; Q7Z5S4; Q8N8B0; B2RPM6
Molecular Weight	96,150 Da
NCBI Official Full Name	semaphorin-4D isoform 1
NCBI Official Synonym Full Names	sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D
NCBI Official Symbol	SEMA4D
NCBI Official Synonym Symbols	CD100; SEMAJ; coll-4; C9orf164; M-sema-G
NCBI Protein Information	semaphorin-4D; A8; GR3; BB18; sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, 4D
UniProt Protein Name	Semaphorin-4D
UniProt Gene Name	SEMA4D
UniProt Synonym Gene Names	C9orf164; CD100; SEMAJ
UniProt Entry Name	SEM4D_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.