

STAT4 recombinant antibody

STAT4 Recombinant Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA278966	HEK293F Cell	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

STAT4; Monoclonal Recombinant Antibody; STAT4 Recombinant Monoclonal Antibody; Signal transducer and activator of transcription 4; STAT4 recombinant antibody

PRODUCT SPECIFICATIONS


Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	26G2
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for STAT4 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human STAT4
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling; Cancer; Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HepG2 cell with AAA278966 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 515-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA278966 diluted at 1:100 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of AAA278966 diluted at 1:100 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for STAT4 recombinant antibody	Carries out a dual signal transduction and activation of transcription. Involved in IL12 signaling.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	345110660
NCBI GeneID	6775
NCBI Accession #	NP_001230764.1
NCBI GenBank Nucleotide #	NM_001243835.1
UniProt Accession #	Q14765; Q96NZ6
Molecular Weight	85,941 Da
NCBI Official Full Name	signal transducer and activator of transcription 4
NCBI Official Synonym Full Names	signal transducer and activator of transcription 4
NCBI Official Symbol	STAT4

NCBI Official Synonym Symbols	SLEB11
NCBI Protein Information	signal transducer and activator of transcription 4
UniProt Protein Name	Signal transducer and activator of transcription 4
UniProt Gene Name	STAT4
UniProt Entry Name	STAT4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/stat4-recombinant-antibody-clone-26g2-immunofluorescence-immunohistochemistry-elisa/278966>