

Mouse anti-Human EIF2AK3 Monoclonal Antibody | anti-PERK antibody

Mouse anti-Homo sapiens (Human) EIF2AK3 Monoclonal Antibody antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279037	Mouse	Human	Immunofluorescence, ELISA


PRODUCT SYNONYMS

EIF2AK3; Monoclonal Antibody; Mouse anti-Homo sapiens (Human) EIF2AK3 Monoclonal Antibody antibody; Phospho-PERK (T981) Monoclonal Antibody; DKFZp781H1925; E2AK3_HUMAN; EC 2.7.11.1; Eif2ak3; Eukaryotic translation initiation factor 2 alpha kinase 3; Eukaryotic translation initiation factor 2-alpha kinase 3; Heme regulated EIF2 alpha kinase; HRI; HsPEK; Pancreatic eIF2 alpha kinase; Pancreatic eIF2-alpha kinase; PEK; PRKR like endoplasmic reticulum kinase; PRKR-like endoplasmic reticulum kinase; WRS; anti-PERK antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	4A3A10
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid; 0.03% Proclin 300, 50% glycerol, 0.01M PBS, pH 7.4.
Applicable Applications for anti-PERK antibody	IF (Immunofluorescence), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Eukaryotic translation initiation factor 2-alpha kinase 3 protein (115-330AA)
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of U87 cells with AAA279037 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).



IF (Immunofluorescence)
Immunofluorescence staining of HeLa cells with AAA279037 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).



IF (Immunofluorescence)
Immunofluorescence staining of A549 cells with AAA279037 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).

NCBI AND UNIPROT INFORMATION

NCBI GI #	134304838
NCBI GeneID	9451
NCBI Accession #	NP_004827.4
NCBI GenBank Nucleotide #	NM_004836.5
UniProt Accession #	Q9NZJ5; O95846; Q53QY0; Q53SB1; A0AVH1; A0AVH2; B2RCU9
Molecular Weight	125,216 Da
NCBI Official Full Name	eukaryotic translation initiation factor 2-alpha kinase 3
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 2-alpha kinase 3
NCBI Official Symbol	EIF2AK3
NCBI Official Synonym Symbols	PEK; WRS; PERK
NCBI Protein Information	eukaryotic translation initiation factor 2-alpha kinase 3; hsPEK; pancreatic EIF2-alpha kinase; PRKR-like endoplasmic reticulum kinase; eukaryotic translation initiation factor 2 alpha kinase 3
UniProt Protein Name	Eukaryotic translation initiation factor 2-alpha kinase 3

UniProt Gene Name	EIF2AK3
UniProt Synonym Gene Names	PEK; PERK; HsPEK
UniProt Entry Name	E2AK3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/eif2ak3-antibody-clone-4a3a10-immunofluorescence-elisa/279037>