

CA1 recombinant antibody

CA1 Recombinant Monoclonal Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279088	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

CA1; Monoclonal Recombinant Antibody; CA1 Recombinant Monoclonal Antibody; Carbonic anhydrase 1 (EC 4.2.1.1)(Carbonate dehydratase I)(Carbonic anhydrase B)(CAB)(Carbonic anhydrase I)(CA-I); CA1 recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	26E3
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for CA1 recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human CA1
Conjugation	Non-conjugated
Research Area	Cardiovascular; Cell biology
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



IHC (Immunohiostchemistry)
IHC image of AAA279088 diluted at 1:50 and staining in paraffin-embedded human spleen tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.36% DAB.

FCM/FACS (Flow Cytometry)
Overlay Peak curve showing MCF-7 cells stained with AAA279088 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4°. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4°. Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.



WB (Western Blot)
Western BlotPositive WB detected in: K562 whole cell lysate,U937 whole cell lysateAll lanes: Ikaros antibody at 1:1000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 28 kDaObserved band size: 28 kDa

NCBI AND UNIPROT INFORMATION

NCBI GI #	192447430
NCBI GeneID	759
NCBI Accession #	NP_001122301.1
NCBI GenBank Nucleotide #	NM_001128829.2
UniProt Accession #	P00915
Molecular Weight	28,870 Da
NCBI Official Full Name	carbonic anhydrase 1
NCBI Official Synonym Full Names	carbonic anhydrase I
NCBI Official Symbol	CA1
NCBI Official Synonym Symbols	CAB; CA-I; Car1
NCBI Protein Information	carbonic anhydrase 1; carbonic anhydrase B; carbonic dehydratase; carbonate dehydratase I

UniProt Protein Name	Carbonic anhydrase 1
UniProt Gene Name	CA1
UniProt Synonym Gene Names	CAB; CA-I
UniProt Entry Name	CAH1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/ca1-recombinant-antibody-clone-26e3-flow-cytometry-immunohistochemistry-western-blot-elisa/279088>