

ASS1 recombinant antibody

ASS1 Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279109	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

ASS1; Monoclonal Recombinant Antibody; ASS1 Recombinant Monoclonal Antibody; Argininosuccinate synthase (EC 6.3.4.5)(Citrulline--aspartate ligase); ASS; ASS1 recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	18G11
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for ASS1 recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human ASS1
Conjugation	Non-conjugated
Research Area	Cancer; Metabolism; Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.




FCM/FACS (Flow Cytometry)

Overlay Peak curve showing Jurkat cells stained with AAA279109 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?.Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.



IHC (Immunohistochemistry)

IHC image of AAA279109 diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.56% DAB.



IF (Immunofluorescence)

Immunofluorescence staining of HeLa with AAA279109 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 516-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

NCBI AND UNIPROT INFORMATION

NCBI GI #	53759107
NCBI GeneID	445
NCBI Accession #	NP_000041.2
NCBI GenBank Nucleotide #	NM_000050.4
UniProt Accession #	P00966; Q6LDL2; Q86UZ0; Q96GT4
Molecular Weight	46,530 Da
NCBI Official Full Name	argininosuccinate synthase
NCBI Official Synonym Full Names	argininosuccinate synthase 1
NCBI Official Symbol	ASS1
NCBI Official Synonym Symbols	ASS; CTLN1
NCBI Protein Information	argininosuccinate synthase; argininosuccinate synthetase 1; citrulline--aspartate ligase; citrulline-aspartate ligase

UniProt Protein Name	Argininosuccinate synthase
UniProt Gene Name	ASS1
UniProt Synonym Gene Names	ASS
UniProt Entry Name	ASSY_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/ass1-recombinant-antibody-clone-18g11-flow-cytometry-immunofluorescence-immunohistochemistry-elisa/279109>