

Rabbit PSMC6 Polyclonal Antibody | anti-PSMC6 antibody

PSMC6 Polyclonal Antibody

CATALOG # AAA279971	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PSMC6; Polyclonal Antibody; PSMC6 Polyclonal Antibody; P44; p42; SUG2; CADP44; anti-PSMC6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	389
Applicable Applications for anti-PSMC6 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human PSMC6
Calculated Molecular Weight	44kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney using PSMC6 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PSMC6 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMC6 antibody	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. Pseudogenes have been identified on chromosomes 8 and 12.
Product Categories/Family for anti-PSMC6 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	51702772
NCBI GeneID	5706
UniProt Accession #	P62333; P49719; Q6IBU3; Q92524; B2R975
Molecular Weight	389
NCBI Official Full Name	26S protease regulatory subunit 10B
NCBI Official Synonym Full Names	proteasome (prosome, macropain) 26S subunit, ATPase, 6
NCBI Official Symbol	PSMC6
NCBI Official Synonym Symbols	P44; p42; SUG2; CADP44; HEL-S-73
NCBI Protein Information	26S protease regulatory subunit 10B; proteasome subunit p42; proteasome 26S subunit ATPase 6; conserved ATPase domain protein 44; epididymis secretory protein Li 73; 26S protease regulatory subunit S10B; 26S proteasome AAA-ATPase subunit RPT4
UniProt Protein Name	26S protease regulatory subunit 10B
UniProt Gene Name	PSMC6
UniProt Synonym Gene Names	SUG2
UniProt Entry Name	PRS10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/psmc6-antibody-immunohistochemistry-western-blot-elisa/279971>