

Rabbit GPD1 Polyclonal Antibody | anti-GPD1 antibody

GPD1 Polyclonal Antibody

CATALOG # AAA279983	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

GPD1; Polyclonal Antibody; GPD1 Polyclonal Antibody; GPD-C; HTGTI; GPDH-C; anti-GPD1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	349
Applicable Applications for anti-GPD1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human GPD1
Calculated Molecular Weight	37kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human breast adipocytes using GPD1 Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat heart using GPD1 Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat heart using GPD1 Antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using GPD1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-GPD1 antibody	This gene encodes a member of the NAD-dependent glycerol-3-phosphate dehydrogenase family. The encoded protein plays a critical role in carbohydrate and lipid metabolism by catalyzing the reversible conversion of dihydroxyacetone phosphate (DHAP) and reduced nicotinic adenine dinucleotide (NADH) to glycerol-3-phosphate (G3P) and NAD ⁺ . The encoded cytosolic protein and mitochondrial glycerol-3-phosphate dehydrogenase also form a glycerol phosphate shuttle that facilitates the transfer of reducing equivalents from the cytosol to mitochondria. Mutations in this gene are a cause of transient infantile hypertriglyceridemia. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
Product Categories/Family for anti-GPD1 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	134047785
NCBI GeneID	2819
UniProt Accession #	P21695; Q8N1B0; F8W1L5
Molecular Weight	349
NCBI Official Full Name	Glycerol-3-phosphate dehydrogenase
NCBI Official Synonym Full Names	glycerol-3-phosphate dehydrogenase 1 (soluble)
NCBI Official Symbol	GPD1
NCBI Official Synonym Symbols	GPD-C; HTGTI; GPDH-C
NCBI Protein Information	glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic; glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic; glycerophosphate dehydrogenase; glycerol-3-phosphate dehydrogenase [NAD+], cytoplasmic
UniProt Protein Name	Glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic
UniProt Gene Name	GPD1
UniProt Synonym Gene Names	GPD-C; GPDH-C
UniProt Entry Name	GPDA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

