

Rabbit GSTA2 Polyclonal Antibody | anti-GSTA2 antibody

GSTA2 Polyclonal Antibody

CATALOG # AAA280380	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

GSTA2; Polyclonal Antibody; GSTA2 Polyclonal Antibody; anti-GSTA2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	222
Applicable Applications for anti-GSTA2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human GSTA2
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human prostate cancer using GSTA2 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using GSTA2 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-GSTA2 antibody	Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes function in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding these enzymes are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of some drugs. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase belonging to the alpha class. The alpha class genes, located in a cluster mapped to chromosome 6, are the most abundantly expressed glutathione S-transferases in liver. In addition to metabolizing bilirubin and certain anti-cancer drugs in the liver, the alpha class of these enzymes exhibit glutathione peroxidase activity thereby protecting the cells from reactive oxygen species and the products of peroxidation.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	215276987
NCBI GeneID	2939
NCBI Accession #	NP_000837.3
NCBI GenBank Nucleotide #	NM_000846.4
UniProt Accession #	P09210; Q12759; Q16491; Q9NTY6
Molecular Weight	26kDa
NCBI Official Full Name	glutathione S-transferase A2
NCBI Official Synonym Full Names	glutathione S-transferase alpha 2
NCBI Official Symbol	GSTA2
NCBI Official Synonym Symbols	GST2; GTA2; GTH2; GSTA2-2
NCBI Protein Information	glutathione S-transferase A2
UniProt Protein Name	Glutathione S-transferase A2
UniProt Gene Name	GSTA2
UniProt Synonym Gene Names	GST2
UniProt Entry Name	GSTA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/gsta2-antibody-immunohistochemistry-western-blot-elisa/280380>