

Rabbit PLCG1 Polyclonal Antibody | anti-PLCG1 antibody

PLCG1 Polyclonal Antibody

CATALOG # AAA280393	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PLCG1; Polyclonal Antibody; PLCG1 Polyclonal Antibody; anti-PLCG1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	1291
Applicable Applications for anti-PLCG1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Route	Synthetic Peptide
Immunogen	A synthetic peptide of human PLCG1
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using PLCG1 Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using PLCG1 Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat brain using PLCG1 Antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PLCG1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-PLCG1 antibody

The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene.

NCBI AND UNIPROT INFORMATION

NCBI GI #	33598948
NCBI GeneID	5335
NCBI Accession #	NP_002651.2
NCBI GenBank Nucleotide #	NM_002660.2
UniProt Accession #	P19174; Q2V575; B7ZLY7; B9EGH4; E1P5W4
Molecular Weight	149kDa
NCBI Official Full Name	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1 isoform a
NCBI Official Synonym Full Names	phospholipase C, gamma 1
NCBI Official Symbol	PLCG1
NCBI Official Synonym Symbols	PLC1; NCKAP3; PLC-II; PLC148; PLCgamma1
NCBI Protein Information	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1
UniProt Protein Name	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1
UniProt Gene Name	PLCG1
UniProt Synonym Gene Names	PLC1; PLC-II; PLC-gamma-1
UniProt Entry Name	PLCG1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

