

Rabbit ADAM15 Polyclonal Antibody | anti-ADAM15 antibody

ADAM15 Polyclonal Antibody

CATALOG # AAA280394	HOST Rabbit	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	---


PRODUCT SYNONYMS

ADAM15; Polyclonal Antibody; ADAM15 Polyclonal Antibody; anti-ADAM15 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence Length	824
Applicable Applications for anti-ADAM15 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant protein of human ADAM15
Species Reactivity/Cross-Reactivity	Human, Mouse, Rat
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cells using ADAM15 Polyclonal Antibody (AAA280394) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using ADAM15 antibody at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-ADAM15 antibody	The protein encoded by this gene is a member of the ADAM (a disintegrin and metalloproteinase) protein family. ADAM family members are type I transmembrane glycoproteins known to be involved in cell adhesion and proteolytic ectodomain processing of cytokines and adhesion molecules. This protein contains multiple functional domains including a zinc-binding metalloprotease domain, a disintegrin-like domain, as well as a EGF-like domain. Through its disintegrin-like domain, this protein specifically interacts with the integrin beta chain, beta 3. It also interacts with Src family protein-tyrosine kinases in a phosphorylation-dependent manner, suggesting that this protein may function in cell-cell adhesion as well as in cellular signaling. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	387849014
NCBI GeneID	8751
NCBI Accession #	NP_001248393.1
NCBI GenBank Nucleotide #	NM_001261464.1
UniProt Accession #	Q13444; Q13493; Q53XQ0; Q5SR68; Q5SR69; Q6R267; Q71S61; Q71S62; B3KQU5; B4DLB5; B4DMH8; E9PN65
Molecular Weight	Observed MW: 93kDa Calculated MW: 55kDa/68kDa/83-92kDa
NCBI Official Full Name	disintegrin and metalloproteinase domain-containing protein 15 isoform 7 preproprotein
NCBI Official Synonym Full Names	ADAM metalloproteinase domain 15
NCBI Official Symbol	ADAM15
NCBI Official Synonym Symbols	MDC15
NCBI Protein Information	disintegrin and metalloproteinase domain-containing protein 15
UniProt Protein Name	Disintegrin and metalloproteinase domain-containing protein 15
UniProt Gene Name	ADAM15
UniProt Synonym Gene Names	MDC15; ADAM 15; MDC-15
UniProt Entry Name	ADA15_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/adam15-antibody-immunofluorescence-western-blot-elisa/280394>