

Rabbit IL5 Polyclonal Antibody | anti-IL5 antibody

IL5 Polyclonal Antibody

CATALOG # AAA280445	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

IL5; Polyclonal Antibody; IL5 Polyclonal Antibody; anti-IL5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	134
Applicable Applications for anti-IL5 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence Corresponding to amino acids 20-134 of human IL5.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using IL5 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver using IL5 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using IL5 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 15s.

ADDITIONAL INFORMATION

Related Product Information for anti-IL5 antibody

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2. This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504671
NCBI GeneID	3567
NCBI Accession #	NP_000870.1
NCBI GenBank Nucleotide #	NM_000879.2
UniProt Accession #	P05113; Q13840
Molecular Weight	15kDa
NCBI Official Full Name	interleukin-5
NCBI Official Synonym Full Names	interleukin 5
NCBI Official Symbol	IL5
NCBI Official Synonym Symbols	EDF; TRF; IL-5
NCBI Protein Information	interleukin-5
UniProt Protein Name	Interleukin-5
UniProt Gene Name	IL5
UniProt Synonym Gene Names	IL-5; TRF
UniProt Entry Name	IL5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

