

Rabbit anti-Mouse, Rat HRG Polyclonal Antibody | anti-HRG antibody

HRG Polyclonal Antibody

CATALOG # AAA280528	HOST Rabbit	REACTIVITY Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	---------------------------------	-------------------------------------


PRODUCT SYNONYMS

HRG; Polyclonal Antibody; HRG Polyclonal Antibody; anti-HRG antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	525
Applicable Applications for anti-HRG antibody	WB (Western Blot)
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human HRG
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse stomach using HRG antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using HRG antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat lung using HRG antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of mouse pancreas, using HRG antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

ADDITIONAL INFORMATION

Related Product Information for anti-HRG antibody

This histidine-rich glycoprotein contains two cystatin-like domains and is located in plasma and platelets. The physiological function has not been determined but it is known that the protein binds heme, dyes and divalent metal ions. The encoded protein also has a peptide that displays antimicrobial activity against *C. albicans*, *E. coli*, *S. aureus*, *P. aeruginosa*, and *E. faecalis*. It can inhibit rosette formation and interacts with heparin, thrombospondin and plasminogen. Two of the protein's effects, the inhibition of fibrinolysis and the reduction of inhibition of coagulation, indicate a potential prothrombotic effect. Mutations in this gene lead to thrombophilia due to abnormal histidine-rich glycoprotein levels.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504489
NCBI GeneID	3273
NCBI Accession #	NP_000403.1
NCBI GenBank Nucleotide #	NM_000412.4
UniProt Accession #	P04196; B9EK35; D3DNU7
Molecular Weight	60kDa
NCBI Official Full Name	histidine-rich glycoprotein
NCBI Official Synonym Full Names	histidine-rich glycoprotein
NCBI Official Symbol	HRG
NCBI Official Synonym Symbols	HPRG; HRGP; THPH11
NCBI Protein Information	histidine-rich glycoprotein
UniProt Protein Name	Histidine-rich glycoprotein
UniProt Gene Name	HRG
UniProt Synonym Gene Names	HPRG
UniProt Entry Name	HRG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

