

Rabbit PCSK9 Polyclonal Antibody | anti-PCSK9 antibody

PCSK9 Polyclonal Antibody

CATALOG # AAA280590	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

PCSK9; Polyclonal Antibody; PCSK9 Polyclonal Antibody; anti-PCSK9 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	692
Applicable Applications for anti-PCSK9 antibody	WB (Western Blot)
Route	Synthetic Peptide
Immunogen	A synthetic peptide of human PCSK9
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse liver using PCSK9 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using PCSK9 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat kidney using PCSK9 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of SW480 cells, using PCSK9 Antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-PCSK9 antibody

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an autocatalytic processing event with its prosegment in the ER and is constitutively secreted as an inactive protease into the extracellular matrix and trans-Golgi network. It is expressed in liver, intestine and kidney tissues and escorts specific receptors for lysosomal degradation. It plays a role in cholesterol and fatty acid metabolism. Mutations in this gene have been associated with autosomal dominant familial hypercholesterolemia. Alternative splicing results in multiple transcript variants.

NCBI AND UNIPROT INFORMATION

NCBI GI #	31317307
NCBI GeneID	255738
NCBI Accession #	NP_777596.2
NCBI GenBank Nucleotide #	NM_174936.3
UniProt Accession #	Q8NBP7; Q5PSM5; Q5SZQ2; A8T640; C0JYY9
Molecular Weight	74kDa
NCBI Official Full Name	proprotein convertase subtilisin/kexin type 9 preproprotein
NCBI Official Synonym Full Names	proprotein convertase subtilisin/kexin type 9
NCBI Official Symbol	PCSK9
NCBI Official Synonym Symbols	FH3; PC9; NARC1; LDLCQ1; NARC-1; HCHOLA3
NCBI Protein Information	proprotein convertase subtilisin/kexin type 9
UniProt Protein Name	Proprotein convertase subtilisin/kexin type 9
UniProt Gene Name	PCSK9
UniProt Synonym Gene Names	NARC1; NARC-1; PC9
UniProt Entry Name	PCSK9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

