

Rabbit CCL11 Polyclonal Antibody | anti-CCL11 antibody

CCL11 Polyclonal Antibody

CATALOG # AAA280624	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

CCL11; Polyclonal Antibody; CCL11 Polyclonal Antibody; anti-CCL11 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	97
Applicable Applications for anti-CCL11 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human CCL11
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of A549 cells using CCL11 antibody. Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using CCL11 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat brain using CCL11 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat lung using CCL11 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CCL11 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-CCL11 antibody

This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, displays chemotactic activity for eosinophils, but not mononuclear cells or neutrophils. This eosinophil-specific chemokine is thought to be involved in eosinophilic inflammatory diseases such as atopic dermatitis, allergic rhinitis, asthma and parasitic infections.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506827
NCBI GeneID	6356
NCBI Accession #	NP_002977.1
NCBI GenBank Nucleotide #	NM_002986.2
UniProt Accession #	P51671; P50877; Q92490; Q92491
Molecular Weight	11kDa
NCBI Official Full Name	eotaxin
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 11

NCBI Official Symbol	CCL11
NCBI Official Synonym Symbols	SCYA11
NCBI Protein Information	eotaxin
UniProt Protein Name	Eotaxin
UniProt Gene Name	CCL11
UniProt Synonym Gene Names	SCYA11
UniProt Entry Name	CCL11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ccl11-antibody-immunofluorescence-western-blot-elisa/280624>