

Rabbit MRC1 Polyclonal Antibody | anti-MRC1 antibody

MRC1 Polyclonal Antibody

CATALOG # AAA280670	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

MRC1; Polyclonal Antibody; MRC1 Polyclonal Antibody; MRC1;CD206;CLEC13D;CLEC13DL;MMR;MRC1L1;bA541I19.1;hMR; anti-MRC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Applicable Applications for anti-MRC1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 610-860 of human MRC1 (NP_002429.1).
Preparation and Storage	Store at -20°C. Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human lung using MRC1 antibody (AAA280670) at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Mouse prostate using MRC1 antibody (AAA280670) at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Rat lung using MRC1 antibody (AAA280670) at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using MRC1 antibody (AAA280670) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-MRC1 antibody	The recognition of complex carbohydrate structures on glycoproteins is an important part of several biological processes, including cell-cell recognition, serum glycoprotein turnover, and neutralization of pathogens. The protein encoded by this gene is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. The protein has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment.
Product Categories/Family for anti-MRC1 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505245
NCBI GeneID	4360
NCBI Accession #	NP_002429.1
NCBI GenBank Nucleotide #	NM_002438.3
UniProt Accession #	P22897; Q5VSJ2; Q5VSK2; A5PKW3
Molecular Weight	Observed MW: 190kDa Calculated MW: 56kDa/166kDa
NCBI Official Full Name	macrophage mannose receptor 1
NCBI Official Synonym Full Names	mannose receptor, C type 1
NCBI Official Symbol	MRC1
NCBI Official Synonym Symbols	MMR; hMR; CD206; MRC1L1; CLEC13D; CLEC13DL; bA541I19.1
NCBI Protein Information	macrophage mannose receptor 1
UniProt Protein Name	Macrophage mannose receptor 1
UniProt Gene Name	MRC1
UniProt Synonym Gene Names	CLEC13D; CLEC13DL; MRC1L1; MMR; hMR
UniProt Entry Name	MRC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mrc1-antibody-immunohistochemistry-western-blot-elisa/280670>