

Rabbit MAPK1 Polyclonal Antibody | anti-MAPK1 antibody

MAPK1 Polyclonal Antibody

CATALOG # AAA280800	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

MAPK1; Polyclonal Antibody; MAPK1 Polyclonal Antibody; ERK; ERK-2; ERK2; ERT1; MAPK2; P42MAPK; PRKM1; PRKM2; p38; p40; p41; p41mapk; p42-MAPK; anti-MAPK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.
Sequence Length	360
Applicable Applications for anti-MAPK1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human MAPK1
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using MAPK1 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using MAPK1 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human prostate using MAPK1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver injury using MAPK1 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using MAPK1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK1 antibody

This gene encodes a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. One study also suggests that this protein acts as a transcriptional repressor independent of its kinase activity. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Two alternatively spliced transcript variants encoding the same protein, but differing in the UTRs, have been reported for this gene.

Product Categories/Family for anti-MAPK1 antibody

[Primary antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	119554
NCBI GeneID	5594
UniProt Accession #	P28482; A8CZ64
Molecular Weight	41kDa
NCBI Official Full Name	Mitogen-activated protein kinase 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase 1

NCBI Official Symbol	MAPK1
NCBI Official Synonym Symbols	ERK; p38; p40; p41; ERK2; ERT1; ERK-2; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk; p42-MAPK
NCBI Protein Information	mitogen-activated protein kinase 1
UniProt Protein Name	Mitogen-activated protein kinase 1
UniProt Gene Name	MAPK1
UniProt Synonym Gene Names	ERK2; PRKM1; PRKM2; MAP kinase 1; MAPK 1; ERK-2; p42-MAPK; MAP kinase 2; MAPK 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/mapk1-antibody-immunohistochemistry-western-blot-elisa/280800>