

Rabbit anti-Human, Mouse STAT3 Polyclonal Antibody | anti-STAT3 antibody

STAT3 Polyclonal Antibody

CATALOG # AAA280804	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	-----------------------------------	---


PRODUCT SYNONYMS

STAT3; Polyclonal Antibody; STAT3 Polyclonal Antibody; ADMIO; ADMIO1; APRE; HIES; anti-STAT3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	770
Applicable Applications for anti-STAT3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human STAT3
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human gastric cancer using STAT3 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon carcinoma using STAT3 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human lung cancer using STAT3 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human tonsil using STAT3 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using STAT3 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-STAT3 antibody

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Product Categories/Family for anti-STAT3 antibody

[Primary antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	48429227
NCBI GeneID	6774
UniProt Accession #	P40763 ; O14916 ; Q9BW54 ; A8K7B8 ; K7ENL3
Molecular Weight	88kDa
NCBI Official Full Name	Signal transducer and activator of transcription 3

NCBI Official Synonym Full Names	signal transducer and activator of transcription 3
NCBI Official Symbol	STAT3
NCBI Official Synonym Symbols	APRF; HIES; ADMIO; ADMIO1
NCBI Protein Information	signal transducer and activator of transcription 3
UniProt Protein Name	Signal transducer and activator of transcription 3
UniProt Gene Name	STAT3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/stat3-antibody-immunohistochemistry-western-blot-elisa/280804>