

Rabbit anti-Human, Mouse NPC2 Polyclonal Antibody | anti-NPC2 antibody

NPC2 Polyclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28086	Rabbit	Human, Mouse	Western Blot, Immunohistochemistry, Immunofluorescence

PRODUCT SYNONYMS


NPC2; Polyclonal Antibody; NPC2 Polyclonal Antibody; HE1; EDDM1; anti-NPC2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	151
Applicable Applications for anti-NPC2 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Application Notes	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF: 1:50 - 1:200
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human NPC2
Calculated Molecular Weight	17kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse lung using NPC2 Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human esophagus using NPC2 Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human prostate using NPC2 Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human colon using NPC2 Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human liver injury using NPC2 Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human lung injury using NPC2 Antibody at dilution of 1:100 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-NPC2 antibody	This gene encodes a protein containing a lipid recognition domain. The encoded protein may function in regulating the transport of cholesterol through the late endosomal/lysosomal system. Mutations in this gene have been associated with Niemann-Pick disease, type C2 and frontal lobe atrophy.
Product Categories/Family for anti-NPC2 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	48429027
NCBI GeneID	10577
UniProt Accession #	P61916; Q15668; Q29413; B4DQV7
Molecular Weight	151
NCBI Official Full Name	Epididymal secretory protein E1
NCBI Official Synonym Full Names	Niemann-Pick disease, type C2
NCBI Official Symbol	NPC2
NCBI Official Synonym Symbols	HE1; EDDM1
NCBI Protein Information	epididymal secretory protein E1; epididymal protein 1; tissue-specific secretory protein; human epididymis-specific protein 1; Niemann-Pick disease type C2 protein
UniProt Protein Name	Epididymal secretory protein E1
UniProt Gene Name	NPC2
UniProt Synonym Gene Names	HE1; He1
UniProt Entry Name	NPC2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/npc2-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/28086>