

Rabbit FLT3 Polyclonal Antibody | anti-FLT3 antibody

FLT3 Polyclonal Antibody

CATALOG # AAA280876	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

FLT3; Polyclonal Antibody; FLT3 Polyclonal Antibody; CD135; FLK-2; FLK2; STK1; anti-FLT3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	993
Applicable Applications for anti-FLT3 antibody	WB (Western Blot)
Immunogen	A synthetic peptide of human FLT3
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver cancer using FLT3 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat pancreas using FLT3 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using FLT3 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-FLT3 antibody

This gene encodes a class III receptor tyrosine kinase that regulates hematopoiesis. This receptor is activated by binding of the fms-related tyrosine kinase 3 ligand to the extracellular domain, which induces homodimer formation in the plasma membrane leading to autophosphorylation of the receptor. The activated receptor kinase subsequently phosphorylates and activates multiple cytoplasmic effector molecules in pathways involved in apoptosis, proliferation, and differentiation of hematopoietic cells in bone marrow. Mutations that result in the constitutive activation of this receptor result in acute myeloid leukemia and acute lymphoblastic leukemia.

Product Categories/Family for anti-FLT3 antibody

Primary antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	156630887
NCBI GeneID	2322
UniProt Accession #	P36888; Q13414; A0AVG9; B7ZLT7; B7ZLT8; F5H0A0
Molecular Weight	113kDa
NCBI Official Full Name	Receptor-type tyrosine-protein kinase FLT3
NCBI Official Synonym Full Names	fms related tyrosine kinase 3
NCBI Official Symbol	FLT3
NCBI Official Synonym Symbols	FLK2; STK1; CD135; FLK-2
NCBI Protein Information	receptor-type tyrosine-protein kinase FLT3
UniProt Protein Name	Receptor-type tyrosine-protein kinase FLT3
UniProt Gene Name	FLT3
UniProt Synonym Gene Names	CD135; FLK2; STK1; FLK-2; FLT-3; STK-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/flt3-antibody-western-blot-elisa/280876>