

Rabbit anti-Human, Mouse XIAP Polyclonal Antibody | anti-XIAP antibody

XIAP Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA280996	Rabbit	Human, Mouse	Immunofluorescence, Western Blot


PRODUCT SYNONYMS

XIAP; Polyclonal Antibody; XIAP Polyclonal Antibody; API3; BIRC4; hIAP-3; hIAP3; IAP-3; ILP1; MIHA; XLP2; anti-XIAP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence	MTFNSFEGSK TCVPADINKE EEFVEEFNRL KTFANFPSGS PVSASTLARA GFLYTGEDT VRCFSCHAAV DRWQYGDSAV GRHRKVSPNC RFINGFYLEN SATQSTNSGI QNGQYKVENY LGSRDHFALD RPSETHADYL LRTGQVVDIS DTIYPRNPAM YSEEARLKSF QNWPDYAHLT PRELASAGLY YTGIGDQVQC
Sequence Length	497
Applicable Applications for anti-XIAP antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant protein of human XIAP
Immunogen Species	Human
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Cellular Location	Cytoplasm, Nucleus
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IF (Immunofluorescence)
Immunofluorescence analysis of HeLa cells using XIAP antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using XIAP antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-XIAP antibody	This gene encodes a protein that belongs to a family of apoptotic suppressor proteins. Members of this family share a conserved motif termed, baculovirus IAP repeat, which is necessary for their anti-apoptotic function. This protein functions through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2 and inhibits apoptosis induced by menadione, a potent inducer of free radicals, and interleukin 1-beta converting enzyme. This protein also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. Mutations in this gene are the cause of X-linked lymphoproliferative syndrome. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 2 and 11.
Product Categories/Family for anti-XIAP antibody	Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	32528299
NCBI GeneID	331
NCBI Accession #	NP_001158.2
NCBI GenBank Nucleotide #	NM_001167.3
UniProt Accession #	P98170; Q9NQ14; D3DTF2
Molecular Weight	Calculated: 56kDa Observed: 51kDa
NCBI Official Full Name	E3 ubiquitin-protein ligase XIAP
NCBI Official Synonym Full Names	X-linked inhibitor of apoptosis
NCBI Official Symbol	XIAP
NCBI Official Synonym Symbols	API3; ILP1; MIHA; XLP2; BIRC4; IAP-3; hIAP3; hIAP-3
NCBI Protein Information	E3 ubiquitin-protein ligase XIAP
UniProt Protein Name	E3 ubiquitin-protein ligase XIAP
UniProt Gene Name	XIAP
UniProt Synonym Gene Names	API3; BIRC4; IAP3; ILP; hILP; IAP-3; hIAP-3; hIAP3; X-linked IAP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/xiap-antibody-immunofluorescence-western-blot-elisa/280996>