

Rabbit FURIN Polyclonal Antibody | anti-FURIN antibody

FURIN Polyclonal Antibody

CATALOG # AAA281017	HOST Rabbit	APPLICATIONS Immunofluorescence, Western Blot, ELISA
-------------------------------	-----------------------	--


PRODUCT SYNONYMS

FURIN; Polyclonal Antibody; FURIN Polyclonal Antibody; FUR; PACE; PCSK3; SPC1; anti-FURIN antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence Length	794
Applicable Applications for anti-FURIN antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 700-794 of human Furin (NP_002560.1).
Species/CrossReactivity	Human, Mouse, Rat
Preparation and Storage	Store at -20°C. Avoid freeze / thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using Furin antibody (AAA281017) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-FURIN antibody	This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. It encodes a type 1 membrane bound protease that is expressed in many tissues, including neuroendocrine, liver, gut, and brain. The encoded protein undergoes an initial autocatalytic processing event in the ER and then sorts to the trans-Golgi network through endosomes where a second autocatalytic event takes place and the catalytic activity is acquired. The product of this gene is one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. Some of its substrates include parathyroid hormone, transforming growth factor beta 1 precursor, proalbumin, pro-beta-secretase, membrane type-1 matrix metalloproteinase, beta subunit of pro-nerve growth factor and von Willebrand factor. It is also thought to be one of the proteases responsible for the activation of HIV envelope glycoproteins gp160 and gp140 and may play a role in tumor progression. This gene is located in close proximity to family member proprotein convertase subtilisin/kexin type 6 and upstream of the FES oncogene. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-FURIN antibody	Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	577019578
NCBI GeneID	5045
NCBI Accession #	NP_001276752.1
NCBI GenBank Nucleotide #	NM_001289823.1
UniProt Accession #	P09958 ; Q14336 ; Q6LBS3 ; Q9UCZ5
Molecular Weight	Observed MW: 150kDa Calculated MW: 114kDa
NCBI Official Full Name	furin preproprotein
NCBI Official Synonym Full Names	furin, paired basic amino acid cleaving enzyme
NCBI Official Symbol	FURIN
NCBI Official Synonym Symbols	FUR; PACE; SPC1; PCSK3
NCBI Protein Information	furin
UniProt Protein Name	Furin
UniProt Gene Name	FURIN
UniProt Synonym Gene Names	FUR; PACE; PCSK3; PACE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/furin-antibody-immunofluorescence-western-blot-elisa/281017>