

Rabbit anti-Human NR3C1 Polyclonal Antibody | anti-NR3C1 antibody

NR3C1 Polyclonal Antibody

CATALOG # AAA281139	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

NR3C1; Polyclonal Antibody; NR3C1 Polyclonal Antibody; GCCR; GCR; GCRST; GR; GRL; anti-NR3C1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence	MDSKESLTPG REENPSSVLA QERGDVMDFY KTLRGGATVK VSASSPSLAV ASQSDSKQRR LLVDFPKGSV SNAQQPDLSK AVSLSMGLYM GETETKVMGN DLGFPQQGQI SLSSGETDLK LLEESIANLN RSTSVPENPK SSASTAVSAA PTEKEFPKTH SDVSSEQHL KGQTGTNGGN VKLYTTDQST FDILQDLEFS SGSPGKETNE SPWRSDLLID ENCLLSPLAG EDDSFLEGN SNEDCKPLIL PDTKP
Sequence Length	777
Applicable Applications for anti-NR3C1 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant protein of human NR3C1
Immunogen Species	Human
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Cellular Location	Cytoplasm, Mitochondrion, Nucleus, Nucleus
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IF (Immunofluorescence)
Immunofluorescence analysis of U2OS cells using NR3C1 antibody at dilution of 1:100.
Blue: DAPI for nuclear staining.



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using NR3C1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-NR3C1 antibody	This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).
Product Categories/Family for anti-NR3C1 antibody	Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504133
NCBI GeneID	2908
NCBI Accession #	NP_000167.1
NCBI GenBank Nucleotide #	NM_000176.2
UniProt Accession #	P04150 ; P04151 ; Q53EP5 ; Q6N0A4 ; A0ZXF9 ; B0LPG8 ; D3DQF4 ; F5ATB7
Molecular Weight	Calculated: 49-85kDa Observed: 105kDa
NCBI Official Full Name	glucocorticoid receptor isoform alpha
NCBI Official Synonym Full Names	nuclear receptor subfamily 3 group C member 1
NCBI Official Symbol	NR3C1
NCBI Official Synonym Symbols	GR; GCR; GRL; GCCR; GCRST
NCBI Protein Information	glucocorticoid receptor
UniProt Protein Name	Glucocorticoid receptor
UniProt Gene Name	NR3C1
UniProt Synonym Gene Names	GRL; GR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/nr3c1-antibody-immunofluorescence-western-blot-elisa/281139>