

Rabbit APRT Polyclonal Antibody | anti-APRT antibody

APRT Polyclonal Antibody

CATALOG # AAA281174	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

APRT; Polyclonal Antibody; APRT Polyclonal Antibody; AMP; APRTD; anti-APRT antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence	MADSELQLVE QRIRSFDFP TPGVVFRDIS PVLKDPASFR AAIGLLARHL KATHGGRIDY IAGLDSRGFL FGPSLAQELG LGCVLIRKRG KLPGPTLWAS YSLEYGKAEL EIQKDALEPG QRVVVDDLL ATGGTMNAAC ELLGRLQAEV LECVSLVELT SLKGREKLAP VPFFSLLQYE
Sequence Length	180
Applicable Applications for anti-APRT antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human APRT
Immunogen Species	Human
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Cellular Location	Cytoplasm
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using APRT Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human prostate using APRT Antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using APRT antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-APRT antibody	Adenine phosphoribosyltransferase belongs to the purine/pyrimidine phosphoribosyltransferase family. A conserved feature of this gene is the distribution of CpG dinucleotides. This enzyme catalyzes the formation of AMP and inorganic pyrophosphate from adenine and 5-phosphoribosyl-1-pyrophosphate (PRPP). It also produces adenine as a by-product of the polyamine biosynthesis pathway. A homozygous deficiency in this enzyme causes 2,8-dihydroxyadenine urolithiasis. Two transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-APRT antibody	Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502171
NCBI GeneID	353
NCBI Accession #	NP_000476.1
NCBI GenBank Nucleotide #	NM_000485.2
UniProt Accession #	P07741 ; Q3KP55 ; Q68DF9 ; G5E9J2
Molecular Weight	Calculated: 14kDa; 19kDa Observed: 19-22kDa
NCBI Official Full Name	adenine phosphoribosyltransferase isoform a
NCBI Official Synonym Full Names	adenine phosphoribosyltransferase
NCBI Official Symbol	APRT
NCBI Official Synonym Symbols	AMP; APRTD
NCBI Protein Information	adenine phosphoribosyltransferase
UniProt Protein Name	Adenine phosphoribosyltransferase
UniProt Gene Name	APRT
UniProt Synonym Gene Names	APRT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

