

Rabbit PSMA5 Polyclonal Antibody | anti-PSMA5 antibody

PSMA5 Polyclonal Antibody

CATALOG # AAA281185	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PSMA5; Polyclonal Antibody; PSMA5 Polyclonal Antibody; PSC5; ZETA; anti-PSMA5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence	MFLTRSEYDR GVNTFSPEGR LFQVEYAIEA IKLGSTAIGI QTSEGVCLAV EKRTSPLME PSSIEKIVEI DAHIGCAMSG LIADAKTLID KARVETQNHWF TYNETMTVE SVTQAVSNLA LQFGEEDADP GAMS RPF GVA LLFGGVDEKG PQLFHMDPSG TFVQCDARAI GSASEGAQSS LQEVYHKSMT LKEAIKSSLI ILKQVMEEKL NATNIELATV QPGQNFHMT KEELEEVIKD I
Sequence Length	183
Applicable Applications for anti-PSMA5 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human PSMA5
Immunogen Species	Human
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Cellular Location	Cytoplasm, Nucleus
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.


IMAGES



IHC (Immunohistochemisry)
Immunohistochemistry of paraffin-embedded mouse lung using PSMA5 Antibody at dilution of 1:100 (40x lens).



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded rat lung using PSMA5 Antibody at dilution of 1:100 (40x lens).



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using PSMA5 antibody at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMA5 antibody	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been found for this gene.
Product Categories/Family for anti-PSMA5 antibody	Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	315138981
NCBI GeneID	5686
NCBI Accession #	NP_001186701.1
NCBI GenBank Nucleotide #	NM_001199772.1
UniProt Accession #	P28066 ; Q3T1C1 ; Q6IBF7 ; B2R8F6 ; B4E2V4
Molecular Weight	Calculated: 20kDa; 26kDa Observed: 28kDa

NCBI Official Full Name	proteasome subunit alpha type-5 isoform 2
NCBI Official Synonym Full Names	proteasome subunit alpha 5
NCBI Official Symbol	PSMA5
NCBI Official Synonym Symbols	PSC5; ZETA
NCBI Protein Information	proteasome subunit alpha type-5
UniProt Protein Name	Proteasome subunit alpha type-5
UniProt Gene Name	PSMA5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/psma5-antibody-immunohistochemistry-western-blot-elisa/281185>