

Rabbit LMNA-S22 Polyclonal Antibody | anti-LMNA-S22 antibody

Phospho-LMNA-S22 pAb

CATALOG # AAA281283	HOST Rabbit	APPLICATIONS Western Blot
-------------------------------	-----------------------	-------------------------------------


PRODUCT SYNONYMS

LMNA-S22; Polyclonal Antibody; Phospho-LMNA-S22 pAb; CDCD1; CDDC; CMD1A; CMT2B1; EMD2; FPL; FPLD; FPLD2; HGPS; IDC; LDP1; LFP; LGMD1B; LMN1; LMNC; LMNL1; MADA; PRO1; anti-LMNA-S22 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.
Sequence Length	574
Applicable Applications for anti-LMNA-S22 antibody	WB (Western Blot)
Immunogen	A synthetic phosphorylated peptide around S22 of human Lamin A (NP_005563.1).
Species Reactivity / Cross Reactivity	Human, Rat
Preparation and Storage	Store at -20°C. Avoid freeze / thaw cycles.

IMAGES



IP (Immunoprecipitation)

Immunoprecipitation analysis of 200 µg extracts of HeLa cells, using 3 µg Phospho-Lamin A/C-S22 pAb (AAA281283). Western blot was performed from the immunoprecipitate using Phospho-Lamin A/C-S22 pAb (AAA281283) at a dilution of 1:1000. HeLa cells were treated by Paclitaxel (100 nM/mL) at 37°C for 20 hours.



WB (Western Blot)

Western blot analysis of various lysates using Phospho-Lamin A/C-S22 Rabbit pAb (AAA281283) at 1:2000 dilution or Lamin A antibody. HeLa cells were treated by Hydroxyurea (4 mM) at 37°C for 20 hours or treated by Paclitaxel (100 nM/mL) at 37°C for 20 hours. C6 cells were treated by Hydroxyurea (4 mM) at 37°C for 20 hours or treated by Paclitaxel (100 nM) at 37°C for 20 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% BSA. Detection: ECL Enhanced Kit. Exposure time: 1s.

ADDITIONAL INFORMATION

Related Product Information for anti-LMNA-S22 antibody	Background: The protein encoded by this gene is part of the nuclear lamina, a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants. Mutations in this gene lead to several diseases: Emery-Dreifuss muscular dystrophy, familial partial lipodystrophy, limb girdle muscular dystrophy, dilated cardiomyopathy, Charcot-Marie-Tooth disease, and Hutchinson-Gilford progeria syndrome.
Product Categories/Family for anti-LMNA-S22 antibody	Phosphorylated Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	383792150
NCBI GeneID	4000
NCBI Accession #	NP_001244303.1
NCBI GenBank Nucleotide #	NM_001257374.2
UniProt Accession #	P02545; P02546; Q5I6Y4; Q5I6Y6; Q5TCJ2; Q5TCJ3; Q6UYC3; Q969I8; B4DI32; D3DVB0; D6RAQ3; E7EUI9
Molecular Weight	Observed MW: 69kDa, 78kDa Calculated MW: 74kDa
NCBI Official Full Name	lamin isoform D
NCBI Official Synonym Full Names	lamin A/C
NCBI Official Symbol	LMNA
NCBI Official Synonym Symbols	FPL; IDC; LFP; CDDC; EMD2; FPLD; HGPS; LDP1; LMN1; LMNC; MADA; PRO1; CDCD1; CMD1A; FPLD2; LMNL1; CMT2B1; LGMD1B
NCBI Protein Information	lamin
UniProt Protein Name	Prelamin-A/C
UniProt Gene Name	LMNA
UniProt Synonym Gene Names	LMN1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/lmna-s22-antibody-western-blot-elisa/281283>