

Rabbit anti-Human, Mouse GSDMD Polyclonal Antibody | anti-GSDMD antibody

GSDMD (Full Length+C terminal) Rabbit pAb

CATALOG # AAA281772	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Western Blot
-------------------------------	-----------------------	-----------------------------------	-------------------------------------


PRODUCT SYNONYMS

GSDMD; Polyclonal Antibody; GSDMD (Full Length+C terminal) Rabbit pAb; DF5L; DFNA5L; FKSG10; GSDMDC1; anti-GSDMD antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-GSDMD antibody	WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 276-484 of human GSDMD (NP_079012.3).
Positive Samples	PC-3, THP-1, RAW264.7
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using GSDMD (Full Length+C terminal) Rabbit pAb at dilution of 1:250 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of PC-12 cells using GSDMD (Full Length+C terminal) Rabbit pAb at dilution of 1:250 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using GSDMD (Full Length+C terminal) Rabbit pAb at dilution of 1:250 (40x lens). Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of RAW264.7 cells, using GSDMD antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 180s.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using GSDMD antibody at 1:1000 dilution. THP-1 cells were treated by PMA/TPA (80 nM) at 37°C for overnight and LPS (1 ug/ml) at 37°C for 6 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-GSDMD antibody	Background: Gasdermin D is a member of the gasdermin family. Members of this family appear to play a role in regulation of epithelial proliferation. Gasdermin D has been suggested to act as a tumor suppressor. Alternatively spliced transcript variants have been described.
Product Categories/Family for anti-GSDMD antibody	Immunology & Inflammation ; Inflammasome Signaling Pathway

NCBI AND UNIPROT INFORMATION

NCBI GI #	83320070
NCBI GeneID	79792
NCBI Accession #	NP_079012.3
NCBI GenBank Nucleotide #	NM_024736.6
UniProt Accession #	P57764 ; Q96Q98 ; D3DWJ9
Molecular Weight	52,801 Da
NCBI Official Full Name	gasdermin-D

NCBI Official Synonym Full Names	gasdermin D
NCBI Official Symbol	GSDMD
NCBI Official Synonym Symbols	DF5L; DFNA5L; FKSG10; GSDMDC1
NCBI Protein Information	gasdermin-D; gasdermin domain containing 1; gasdermin domain-containing protein 1
UniProt Protein Name	Gasdermin-D
UniProt Gene Name	GSDMD
UniProt Synonym Gene Names	DFNA5L; GSDMDC1
UniProt Entry Name	GSDMD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/gsdmd-antibody-western-blot-elisa/281772>