

Rabbit anti-Human, Rat BRD4 Polyclonal Antibody | anti-BRD4 antibody

[KO Validated] BRD4 Rabbit pAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA282031	Rabbit	Human, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

BRD4; Polyclonal Antibody; [KO Validated] BRD4 Rabbit pAb; CAP; HUNK1; HUNKI; MCAP; BRD4 short isoform; anti-BRD4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-BRD4 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human BRD4
Cellular Location	Chromosome, Nucleus
Positive Samples	HeLa, HeLa-BRD4-FL, HeLa-BRD4-S(a), HeLa-BRD4-S(b)
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of extracts from normal (control) and BRD4 Rabbit pAb knockout (KO) HeLa cells, using BRD4 Rabbit pAb antibody at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 180s.



WB (Western Blot)

Western blot analysis of extracts of HeLa cells, using BRD4 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-BRD4 antibody	Background: The protein encoded by this gene is homologous to the murine protein MCAP, which associates with chromosomes during mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. This gene has been implicated as the chromosome 19 target of translocation t(15;19)(q13;p13.1), which defines an upper respiratory tract carcinoma in young people. Two alternatively spliced transcript variants have been described.
Product Categories/Family for anti-BRD4 antibody	Signal Transduction; Kinase; Serine/threonine kinases

NCBI AND UNIPROT INFORMATION

NCBI GI #	7657218
NCBI GeneID	23476
NCBI Accession #	NP_055114.1
NCBI GenBank Nucleotide #	NM_014299.2
UniProt Accession #	O60885; O60433; Q4G0X8; Q86YS8; Q96PD3
Molecular Weight	152kDa
NCBI Official Full Name	bromodomain-containing protein 4 isoform short
NCBI Official Synonym Full Names	bromodomain containing 4
NCBI Official Symbol	BRD4
NCBI Official Synonym Symbols	CAP; MCAP; HUNK1; HUNKI
NCBI Protein Information	bromodomain-containing protein 4; bromodomain-containing 4; chromosome-associated protein
UniProt Protein Name	Bromodomain-containing protein 4
UniProt Gene Name	BRD4
UniProt Synonym Gene Names	HUNK1
UniProt Entry Name	BRD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/brd4-antibody-immunohistochemistry-western-blot-elisa/282031>