

Rabbit anti-Human CD85j Polyclonal Antibody | anti-CD85j antibody

CD85j Rabbit pAb

CATALOG # AAA282212	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot
-------------------------------	-----------------------	----------------------------	-------------------------------------


PRODUCT SYNONYMS

CD85j; Polyclonal Antibody; CD85j Rabbit pAb; LILRB1; CD85J; ILT-2; ILT2; LIR-1; LIR1; MIR-7; MIR7; PIR-B; PIRB; anti-CD85j antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
Applicable Applications for anti-CD85j antibody	WB (Western Blot)
Positive Samples	Daudi, K-562
Immunogen	Recombinant protein of human CD85j.
Cellular Location	cytoplasm, external side of plasma membrane, extracellular region, plasma membrane
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of Daudi cells using CD85j Rabbit pAb at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of various lysates, using CD85j antibody at 1:1190 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.

ADDITIONAL INFORMATION

Related Product Information for anti-CD85j antibody	This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-CD85j antibody	Immunology & Inflammation; CD markers; Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GeneID	10859
Molecular Weight	Observed MW: 110KDa Calculated MW: 49kDa/70kDa/71kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd85j-antibody-western-blot-elisa/282212>