

Rabbit EIF3H Polyclonal Antibody | anti-EIF3H antibody

EIF3H Polyclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28265	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence, Immunoprecipitation

PRODUCT SYNONYMS


EIF3H; Polyclonal Antibody; EIF3H Polyclonal Antibody; eIF3-gamma; eIF3-p40; EIF3S3; anti-EIF3H antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence	NCFPPQHT E DDADFDEVQY QMEMMRSLRH VNIDHLHVGW YQSTYYGSFV TRALLDSQFS YQHAIEESVV LIYDPIKTAQ GSLSLKAYRL TPKLMEVCKE KDFSPEALKK ANITFEYMF E EVPIVIKNSH LINVLMWELE KKSAVADKHE LLSLASSNHL GKNLQLLMDR VDEMSQDIVK YNTYMRNTSK QQQQKHQYQQ RRQQENMQRQ SRGEPPLPEE DLSKLFKPPQ PPARMDSLI AGQINTYCN I KEFT
Sequence Length	352
Applicable Applications for anti-EIF3H antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation)
Application Notes	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF: 1:50 - 1:200 IP: 1:50 - 1:200
Immunogen	Recombinant protein of human EIF3H
Immunogen Species	Human
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Cellular Location	Cytoplasm
Positive Samples	HL-60, HT-29, NIH/3T3, HeLa, Jurkat, Mouse spleen, Mouse testis, Mouse heart, Rat spleen
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.




IP (Immunoprecipitation)
Immunoprecipitation analysis of 200ug extracts of Jurkat cells using 1ug EIF3H antibody. Western blot was performed from the immunoprecipitate using EIF3H antibody at a dilution of 1:1000.




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse heart using EIF3H Antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse brain using EIF3H Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human prostate using EIF3H Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded rat brain using EIF3H Antibody at dilution of 1:100 (40x lens).



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using EIF3H antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

ADDITIONAL INFORMATION

Product Categories/Family for anti-EIF3H antibody	Polyclonal Antibodies
---	-----------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503515
NCBI GeneID	8667
NCBI Accession #	NP_003747.1
NCBI GenBank Nucleotide #	NM_003756.2
UniProt Accession #	O15372
Molecular Weight	Calculated: 39kDa Observed: 40kDa
NCBI Official Full Name	eukaryotic translation initiation factor 3 subunit H
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 3 subunit H
NCBI Official Symbol	EIF3H
NCBI Official Synonym Symbols	EIF3S3; eIF3-p40; eIF3-gamma
NCBI Protein Information	eukaryotic translation initiation factor 3 subunit H
UniProt Protein Name	Eukaryotic translation initiation factor 3 subunit H
UniProt Gene Name	EIF3H
UniProt Synonym Gene Names	eIF3h

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/eif3h-antibody-western-blot-immunohistochemistry-immunofluorescence-immunoprecipitation-elisa/28265>