

Rabbit anti-Human IL17A Monoclonal Antibody | anti-IL17A antibody

IL17A Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA282908	Rabbit	Human	ELISA, Immunohistochemistry


PRODUCT SYNONYMS

IL17A; Monoclonal Antibody; IL17A Rabbit mAb; IL17; CTLA8; IL-17; ILA17; CTLA-8; IL-17A; anti-IL17A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.
Sequence	GITIPRNP GC P NSEDKNFPR TVMVNLNIHN RNTNTNPKRS SDYYNRSTSP WNLHRNEDPE RYPSVIWEAK CRHLGCINAD GNV DYHMNSV PIQ QEILVLR REPPHCPNSF RLEKILVSVG CTCVTPIVHH VA
Applicable Applications for anti-IL17A antibody	ELISA, IHC (Immunohistochemistry)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 24-155 of human IL17A (NP_002181.1).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded Human tonsil using IL17A Rabbit mAb (AAA282908) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using IL17A Rabbit mAb (AAA282908) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.

ADDITIONAL INFORMATION

Related Product Information for anti-IL17A antibody	This gene is a member of the IL-17 receptor family which includes five members (IL-17RA-E) and the encoded protein is a proinflammatory cytokine produced by activated T cells. IL-17A-mediated downstream pathways induce the production of inflammatory molecules, chemokines, antimicrobial peptides, and remodeling proteins. The encoded protein elicits crucial impacts on host defense, cell trafficking, immune modulation, and tissue repair, with a key role in the induction of innate immune defenses. This cytokine stimulates non-hematopoietic cells and promotes chemokine production thereby attracting myeloid cells to inflammatory sites. This cytokine also regulates the activities of NF-kappaB and mitogen-activated protein kinases and can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). IL-17A plays a pivotal role in various infectious diseases, inflammatory and autoimmune disorders, and cancer. High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The lung damage induced by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL17A.
Product Categories/Family for anti-IL17A antibody	Cancer ; Tumor immunology ; Cell Biology Developmental Biology ; Apoptosis ; Immunology Inflammation ; Cytokines ; Interleukins ; Cell Intrinsic Innate Immunity Signaling Pathway ; Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504651
NCBI GeneID	3605
NCBI Accession #	NP_002181.1
NCBI GenBank Nucleotide #	NM_002190.2
UniProt Accession #	Q16552 ; Q5T2P0
Molecular Weight	Calculated MW: 18kDa
UniProt Protein Name	Interleukin-17A
UniProt Gene Name	IL17A
UniProt Synonym Gene Names	CTLA8; IL17; IL-17; IL-17A; CTLA-8
UniProt Entry Name	IL17_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/il17a-antibody-immunohistochemistry-elisa/282908>