

Rabbit anti-Human PMS2 Monoclonal Antibody | anti-PMS2 antibody

[KO Validated] PMS2 Rabbit mAb

CATALOG # AAA282955	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

PMS2; Monoclonal Antibody; [KO Validated] PMS2 Rabbit mAb; MLH4; PMS-2; PMSL2; HNPCC4; LYNCH4; MMRC4; PMS2CL; S2; anti-PMS2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.
Applicable Applications for anti-PMS2 antibody	ELISA, WB (Western Blot)
Immunogen	Recombinant protein of human PMS2.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of lysates from wild type (WT) and PMS2 knockout (KO) HeLa cells, using [KO Validated] PMS2 Rabbit mAb (AAA282955) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



WB (Western Blot)

Western blot analysis of lysates from Jurkat cells, using [KO Validated] PMS2 Rabbit mAb (AAA282955) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-PMS2 antibody	The protein encoded by this gene is a key component of the mismatch repair system that functions to correct DNA mismatches and small insertions and deletions that can occur during DNA replication and homologous recombination. This protein forms heterodimers with the gene product of the mutL homolog 1 (MLH1) gene to form the MutL-alpha heterodimer. The MutL-alpha heterodimer possesses an endonucleolytic activity that is activated following recognition of mismatches and insertion/deletion loops by the MutS-alpha and MutS-beta heterodimers, and is necessary for removal of the mismatched DNA. There is a DQHA(X)2E(X)4E motif found at the C-terminus of the protein encoded by this gene that forms part of the active site of the nuclease. Mutations in this gene have been associated with hereditary nonpolyposis colorectal cancer (HNPCC; also known as Lynch syndrome) and Turcot syndrome.
Product Categories/Family for anti-PMS2 antibody	Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505913
NCBI GeneID	5395
NCBI Accession #	NP_000526.1
NCBI GenBank Nucleotide #	NM_000535.5
UniProt Accession #	P54278; Q52LH6; Q5FBW9; Q5FBX1; Q5FBX2; Q75MR2; B2R610
Molecular Weight	Calculated MW: 96kDa Observed MW: 110kDa
UniProt Protein Name	Mismatch repair endonuclease PMS2
UniProt Gene Name	PMS2
UniProt Synonym Gene Names	PMSL2
UniProt Entry Name	PMS2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/pms2-antibody-western-blot-elisa/282955>